



Anti-tubercular activity of some six membered heterocycle compounds

Mohammad Asif

GRD (PG) Institute of Management and Technology, Dehradun, 248009, India

*Corresponding author's E. mail: asif321@gmail.com

ARTICLE INFO

Article type:

Review article

Article history:

Received February 2014

Accepted May 2015

July 2015 Issue

Keywords:

Anti-tubercular agents

Heterocyclic compounds

Mycobacterium tuberculosis

Drug resistance

ABSTRACT

The effectiveness in TB treatment is difficult because the structural composition of the mycobacterial cell wall is very complicated, which makes many drugs ineffective. Tuberculosis is still one of the most imperative infectious disease worldwide because its important reason is drug resistant, persistent or latent tuberculosis and synergism with HIV. Furthermore no any new chemical entity has come. The recently application of modern drug design promise to bring significant development in the fight against TB. In present review we discussed brief introduction of tuberculosis.

© 2015 International Scientific Organization: All rights reserved.

Capsule Summary: Anti-tubercular activity of six membered heterocycle compounds is reviewed.

Cite This Article As: Asif, M., 2015. Anti-tubercular activity of some six membered heterocycle compounds. Chemistry International 1(3) 134-163.

INTRODUCTION

Tuberculosis (TB) is an infectious disease caused by *Mycobacterium tuberculosis* (Mtb); it is the world's one of common cause of death (Ducati et al., 2006). Aproxymate 2 million people die every year and more than 9 million are getting infected. The present therapy is directly observed treatment short-course (DOTS) and DOTS-Plus (DOTS plus Second-line TB drugs) for Tb and resistant-TB (Loddenkemper et al., 2002; Perri and Bonora, 2004). The TB has been spreading at a steady rate over the last decade (Bishai and Chaisson, 1997) and the recovery in TB is alarming due to the development of pathogenic synergy with HIV. In the current treatment, the emergence of multi drug resistant TB (MDR-TB) and extensively drug resistant TB (XDR-TB) as a consequence of lengthy treatment, makes patient compliance difficult. These problems are encrease for the development of new anti-TB drug. The MDR-TB strains are resistant to two or more of the five first-line anti-TB drugs (isoniazid, rifampicin, pyrazinamide, ethambutol and

streptomycin) (Bastian and Colebunders, 1999; Barry et al., 2000). MDR-TB takes longer to treat with second-line drugs (DOT-Plus), which are more expensive and have more side-effects. XDR-TB will develop when these second-line drugs are mismanaged and also become ineffective. The development of drug resistance that may becomes an incurable disease (Asif, 2012a; Asif, 2012b) and provides a strong motivation for the development of effective and affordable (Asif, 2015a-g; Ashraf et al., 2015) anti-TB agents. Approximately 9 million people developing active TB every year and 1.7 million deaths annually, TB is far from under control. Human immunodeficiency virus (HIV) infection dramatically increases the risk of developing active tuberculosis and is driving the TB epidemic in Africa. HIV renders tuberculosis more difficult to diagnose (due to higher incidence of sputum negative disease), and treat (due to interactions and side-effects). The increasing spread of multidrug-resistant TB (MDR-TB) and the recalcitrant nature of persistent infections pose additional challenges to treatment with currently available anti-TB drugs. The situation is exacerbated by

the increasing emergence of extensively drug-resistant (XDR) TB. Resistance to at least two main first-line drugs and additionally to three or more of the six classes of second-line drugs makes this form of TB virtually untreatable with available drugs. Although TB can be cured, current treatment is complex and long lasting, involving four drugs for 2 months and two drugs for at least another 4 months. Directly Observed Therapy (DOT), as promoted by the World Health Organisation (WHO) to improve compliance for the difficult and long-lasting regimen, is demanding for patients, labour intensive for health staff and is compromised in settings where health services are poorly accessible. MDR-TB is even more complex and expensive to treat, and in developing countries treatment is limited to a few projects with limited numbers of patients. After decades of standstill in TB drug development, the drug pipeline has begun to fill up during the last 5 years (Asif, 2012a-c).

Established in 2000, the Global Alliance for TB Drug Development (TB Alliance) has played a critical role in changing the TB research and development (R&D) landscape and is associated with approximately half of all compounds in development. The main criteria established by the TB Alliance to select drug candidates for further development are shortening of the current treatment, activity against MDR-TB and lack of interactions with antiretroviral drugs represent. During the last years, increased public awareness of the lack of R&D for neglected diseases has led at least one pharmaceutical company to establish an institute undertaking R&D activities in tuberculosis on a 'noprofit-no-loss' basis. Other companies have engaged in tuberculosis R&D on for-profit basis, and with some success: three of the six anti-TB candidate drugs currently in clinical trials have been developed by for-profit companies. Major advances have been also made in basic research. Modern molecular and genetic tools have become available for *Mtb* (such as targeted mutagenesis, array-based analysis of mutant libraries, techniques for conditional gene silencing, and global gene expression profiling) and this has led to impressive improvements in the knowledge and understanding of the basic biology and physiology of *Mtb* (Asif, 2012a-c).

CURRENTLY USED ANTITUBERCULAR AGENT

Isoniazid, ethionamide, prothionamide and pyrazinamide

Isoniazid (INH) (**1**) a potent anti-TB drug with minimal inhibitory concentration (MIC) is 0.05 µg/mL. It acts on growing cells and not on resting cells. INH is kill mycobacteria by inhibiting the biosynthesis of mycolic acids, critical component of the cell wall. The 2-Ethyl thioisonicotinamide (ethionamide) (**2**) and prothionamide (**3**) have MIC 0.5-2.5 µg/mL. The ethionamide also disturbs mycolic acid synthesis in strains resistant to isoniazid, streptomycin and p-amino salicylic acid.

Pyrazinamide (**4**) an analog of nicotinamide is active at a MIC of 6-60 µg/mL. Resistance to pyrazinamide develops soon when it is used alone. Its mechanism of action is unknown (Gray. 1997; Janin. 2007; Sunduru et al., 2010).

NEW POTENTIAL ANTI-TUBERCULAR AGENTS
















The new potential anti-TB agents hav different chemical entities.

Quinolones

Recently, the quinolone drugs; Ofloxacin, Gatifloxacin, Moxifloxacin and Levofloxacin, are serving as second line drugs for TB. In this concern, many researchers optimized the quinolones and evaluated for their anti-TB potency. In this direction, a series of 1-ethyl- and 1-aryl-6-fluoro-1,4-dihydroquinol-4-one derivatives were evaluated for anti-TB and cytotoxic activities. Of these, once derivatives (**5**) exhibited the preeminent MIC of 1.56 µg/mL against *Mtb* H37Rv (MTB) and also a good selectivity index (SI=>40.06). Further, compound **5** also proved to be a potent anti-TB agent with an EC90 value of 5.75 µg/ml (Sheu et al., 2003). Similarly, a number of fifty-one novel 1-(cyclopropyl/2,4-difluorophenyl/t-butyl)-1,4-dihydro-6-fluoro-7-(sub secondary amino)-4-oxoquinoline-3-carboxylic acids and found a potent anti-TB agent (**6**), which showed MIC of 0.09 µM against MTB and MDR-TB respectively. In the in vivo animal model **6** decreased the mycobacterial load in lung and spleen tissues with 2.53- and 4.88-log10 protections respectively at a dose of 50 mg/kg body weight (Senthilkumar et al., 2009).

A series of pyridobenzoxazine derivatives by replacement of the *N*-methylpiperazinyl group of Levofloxacin with various basic substituents to investigate anti-TB activities. Among the compounds, compound **7**, which was a 2,8-diazabicyclo(4.3.0)nonanyl derivative with relatively low lipophilicity, showed the most potent activity against mycobacterial species: the activity was 4- to 32-fold more potent than that of Levofloxacin. These results suggested that an increase in the lipophilicity of Levofloxacin analogues in part contributed to enhancement of anti-TB activities but that lipophilicity of the compound was not a critical factor affecting the potency (Kawakami et al., 2000). While in the investigation of potency against *M. kansasii* Levofloxacin showed MIC in the range of 0.12-0.25 µg/ml while Moxifloxacin showed the range of MIC=≤0.06-0.12 µg/mL (Alcaide et al., 2004). These results prompted for optimization of other quinolone antibacterials to be investigated as anti-TB agents.

Inspired with the activity profile of quinolones, a series of Lamivudine prodrugs bearing fluoroquinolones (**8**) and evaluated their efficacy against *Mtb* H37Rv. All the compounds exhibited an inhibition of 92-100% at a concentration of 6.25 µg/ml (Sriram

**1****2****3****4****5****6****7****8** R₁= ethyl, cyclopropylR₂=H,OCH₃ R₃=H, CH₃ R₄=H,Cl,F**9****10****11****12a:** R=ethyl **12b:** R=cyclopropyl**13****14****15**

et al., 2005). While in ciprofloxacin derivatives, one compound (**9**) showed in vivo anti-TB activity by reducing the bacterial load in spleen tissue with 0.76-log₁₀ protections and was considered to be moderately active in reducing bacterial count in spleen (Sriram et al., 2005). In continuation, Gatifloxacin derivatives

and found a more potent compound (**10**) in comparison to compound **9**. In the in vivo animal model **10** decreased the bacterial load in lung and spleen tissues with 3.62- and 3.76-log₁₀ protections, respectively (Sriram et al., 2006). With this motivation, he was able to find out a most potent molecule (**210**)

which decreased the bacterial load in lung and spleen tissues with 2.42- and 3.66-log₁₀ protections, respectively, at 25 mg/kg body weight (Sriram et al., 2006). Contrarily, 7-(4-(5-amino-1,3,4-thiadiazole-2-sulfonyl))-1-piperazinyl fluoroquinolonic derivatives (**211a** and **211b**), showed moderate anti-TB activity at MIC of 10 µg/mL compared to isoniazid standard (Talatha et al., 2006).

In another approach, 3-unsubstituted 4-hydroxyquinolin-2(1*H*)-one potency against *Mtb* H37Rv, one compound (**13**) showed moderate activity of MIC 3.125 µg/mL (Arya and Agarwal, 2007). Surprisingly, the series of 1-hydroxy-3-oxo-5,6-dihydro-3*H*-pyrrolo(3,2,1-ij)quinoline-2-carboxylic acid hetaryl amides exhibited excellent activity (MIC=0.39-6.25 µg/mL) in comparison to **13**. The most active compound **14** showed MIC of 0.39 µg/mL against *Mtb* H37Rv (Ukrainets et al., 2007). Whereas, the effect of nitro substitution on quinoline ring, a series of 2-(sub)-3-fluoro/nitro-5,12-dihydro-5-oxobenzothiazolo(3,2-a)quinoline-6-carboxylic acid derivatives were evaluated for in-vitro and in-vivo anti-TB activities against *Mtb* H37Rv (MTB), MDR-TB, and *M. smegmatis*, and also tested for the ability to inhibit the supercoiling activity of DNA gyrase from *M. smegmatis*. Among the thirty-four compounds, 2-(3-(diethylcarbamoyl)piperidin-1-yl)-3-fluoro-5,12-dihydro-5-oxobenzothiazolo(3,2-a)quinoline-6-carboxylic acid (**15**) was found to be the most active compound with MIC of 0.18 and 0.08 µM against MTB and MDR-TB, respectively. In the in-vivo animal model **15** decreased the bacterial load in lung and spleen tissues with 2.78 and 3.12- log₁₀ protections, respectively, at the dose of 50 mg/kg body weight (Dinakaran et al., 2008). In another investigation, 6-nitroquinolone (**16**) was also found to be the most active compound in vitro with MIC of 0.08 and 0.16 µM against MTB and MDR-TB, respectively. In the in vivo animal model **16** decreased the bacterial load in lung and spleen tissues with 2.78 and 4.15-log₁₀ protections, respectively, at the dose of 50 mg/kg body weight (Senthilkumar et al., 2009).

In an effort to increase the potency of quinolones, a series of (1,2,3)Triazolo(4,5-*h*)quinolones were evaluated their anti-TB activity against *Mtb* H37Rv and further 11 clinically isolated strains of *Mtb* endowed with different drug resistance. Among all, compound **17** exhibited best activity against all strains with a MIC of 0.5µg/mL (Carta et al., 2007). Whereas in another series of (1,2,3)Triazolo(4,5-*h*)quinolones, Compounds **18** and **19** exhibited better potency of MIC in the range 0.125-16.0 µg/mL against H37Rv and 11 clinical isolates of MDR-TB. These results showed that (1,2,3)-triazolo(4,5-*h*)quinolones were endowed with an excellent activity against MDR-TB strains with no cytotoxicity (Carta et al., 2008).


In the process of investigating novel quinolones as anti-TB agents, many derivatives of quinolones were screened for their in vitro efficacy against MTB and MDR-TB. The most potent (in vitro) compound of the series was screened for in vivo potency

too. Compound **20** exhibited MIC₉₉ of 0.19 µM and 0.09 µM against MTB and MDR-TB, respectively and decreased the bacterial load in lung and spleen tissues with 1.91 and 2.91-log₁₀ protections, respectively, in the in vivo animal model at a dose of 50 mg/kg body weight (Dinakaran, et al., 2008). Compound **21** decreased the bacterial load in lung and spleen tissues with 2.54 and 2.92-log₁₀ protections (Senthilkumar et al., 2008), while **22** decreased the bacterial load by 30% and 42%, respectively, at a dose of 50 mg/kg body weight (Dinakaran et al., 2008). In an effort to increase the anti-TB potency of quinolones, 1-(cyclopropyl/2,4-difluorophenyl/tert-butyl)-1,4-dihydro-8-methyl-6-nitro-4-oxo-7-(substituted-secondary-amino) quinoline-3-carboxylic acids. The most active compound (**23**) of the series showed MIC of 0.42 µM and 0.09 µM against MTB and MDR-TB respectively (Senthilkumar et al., 2009a). While in another series, 7-(3-(diethylcarbamoyl) piperidin-1-yl)-1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxoquinoline-3-carboxylic acid (**24**) exhibited promising MIC of 0.09 µM against MTB and MDR-TB respectively. In the in vivo animal model **24** also decreased the mycobacterial load in lung and spleen tissues with 2.53- and 4.88-log₁₀ protections respectively at a dose of 50 mg/kg body weight (Senthilkumar et al., 2009b).



With the same motivation, Moxifloxacin and Gatifloxacin derivatives and evaluated against *Mtb* H37Rv (MTB), the most active compound (**25**) exhibited a MIC of 0.31µg/mL (de Almeida et al., 2007). While in the series of Tetracycline incorporated with quinolones, compound **26** was found to be the most active against MTB with a MIC of 0.2 µg/mL and also nontoxic to the CEM cells until 200 µM (Sriram et al., 2007). Thus, developing quinolones as anti-TB agents is a worthy approach.

New isoniazid derivatives with extended spectra of activity, a series of isonicotinoylhydrazones, isonicotinohydrazides and their cyanoborane adduct were tested for their *in-vitro* anti-TB activities. Among these, isonicotinohydrazide (**27**) found to be most active, which was able to kill *Mtb* Erdman strain. Compounds **28** showed a MIC of 0.05, 0.1 µg/mL, respectively against *Mtb* H37Rv and Erdman strains (Maccari et al., 2002) and MIC value is <0.025 to <6.25µg/mL. The most potent compound (**29**) also showed good selectivity index of more than 2000 (Hearn et al., 2009). Two series of 4-thiazolidinone and 2-azitidinone derivatives of INH, compound **30** and **31** having 4-hydroxy-3-methoxyphenyl substituent have shown best activity with a MIC 0.31 µg/mL against *Mtb* H37Rv (Jaju et al., 2009). Benzylsulfanylpyridine-2-carbohydrazides (**32**) showed moderate activity against *Mtb*, non-tuberculous *Mycobacterium*, and MDR-TB with MIC values in a range of 2 to 125µM/L (Herzigova et al., 2009).


The 1,4-dihydropyridines recently been shown to possess anti-TB activity. Dihydropyridine derivative **33** was found to be most potent showing 87% inhibition respectively at a




16

17: R=C₄H₉ 18: R=C₂H₅ 19: R=CH₂CH=CH₂


20




21




22




23




24




25




26




27




28




29



30



31



32

concentration of 12.5µg/mL (Gaveriya et al., 2001). While, presence of imidazole group at 4-position and amide group at

3,5-position (34) increased the activity up to 1 µg/mL against *Mtb*. Compound 35 increase the anti-TB activity (Kharkar et al.,

2002). Compound **36** showed best potency with a MIC 1 $\mu\text{M}/\text{mL}$ against *Mtb* H37Rv, which is equal to that of INH (Khoshneviszadeh et al., 2009). 1,4-dihydropyridine-3,5-dicarboxamide derivatives and most active compound (**37**) of the series showed equal potency similar to that of **36** (Fassihi et al., 2009).

Rifabutin (RBT) analogues, compound **38** displayed good potency of MIC $<0.013\mu\text{g}/\text{mL}$ against *Mtb* H37Rv, while compound **39** showed potency of MIC 0.08 μM against non-replicating *Mtb* strains (Figueiredo et al., 2009).

The BM 212 (**40**) with very good *in vitro* activity of MIC 0.7 $\mu\text{g}/\text{mL}$ and compound **41** showed potent inhibition of MIC 0.4 $\mu\text{g}/\text{mL}$ against *Mtb* (Biava et al., 2003). While compounds **42** (Biava et al., 2004) and **43** (Biava et al., 2005) showed comparable MIC of 1 $\mu\text{g}/\text{mL}$. Surprisingly, compound **44** has increased the potency with a MIC of 0.25 $\mu\text{g}/\text{mL}$, equal to isoniazid (INH) (Biava et al., 2008). Compound **45** (Biava et al., 2006) showed decreased potency of MIC 0.4 $\mu\text{g}/\text{mL}$ but lowered the toxicity. This made the molecule a promising lead with a protection index of 160, which is greater than INH and streptomycin. The compound 1-(4-fluorophenyl)-2-ethyl-3-(thiomorpholin-4-yl)methyl-5-(4-methylphenyl)-1H-pyrrole (**46**) is particularly active, with a MIC 0.25 $\mu\text{g}/\text{mL}$ (Biava et al., 2009). All the compounds (**41-46**) were also active against resistant *M. tuberculosis* strains. Compound **47** was found to be most potent with a MIC of 0.5 $\mu\text{g}/\text{mL}$ (Ragno et al., 2000).


The anti-TB efficacy of compounds (**48a-e**) was against TB complex and other non-mycobacterial species. The compounds were significantly active against *Mtb* complex. Compound **48a** showed preminent inhibition of MIC 0.006 mg/L against *M. tuberculosis* UT30 (streptomycin resistant at 4 mg/L). Whereas, compound **48b** showed same potency against *Mtb* UT18 and *M. bovis* BCG. Compound **48d** showed the best potency of all, against *M. bovis* BCG with a MIC of 0.0008 mg/L and also showed the same potency against both the *Mtb* UT15 and UT18. Similarly, Compound **39e** showed the same potency against *Mtb* UT18 but shown increased potency of MIC of 0.0004 mg/L against *Mtb* UT15. Compound **48c** showed the best potency against *M. bovis* BCG with a MIC of 0.0015 mg/L. These NFAs have shown MIC in the range of 0.012-0.006 mg/L in broth assay, 0.012-0.0015 mg/L in agar assay and 0.85- 0.17 in low-oxygen recovery assay against *Mtb* H37Rv (Hurdle et al., 2008; Tangallapally et al., 2006; Tangallapally et al., 2007; Tangallapally et al., 2005).

A pyridyl derivative (**49**) has shown potency with a MIC 0.22 μM and was 3 times more active than standard INH and equally active as RIF in *Mtb* H37Rv. In starved *Mtb* H37Rv, it also inhibited with a MIC of 13.9 μM and was found to be 50 times more active than INH and slightly more active than RIF (Sriram et al., 2009). A series of 5-aryl-1-isonicotinoyl-3-(pyridin-2-yl)-4,5-dihydro-1H-pyrazole derivatives have anti-TB


activity toward *Mtb* H37Rv and *Mtb* H4. Compounds **50a-e** has shown equal potency against above strains with a MIC of 8 $\mu\text{g}/\text{mL}$ (Mamolo et al., 2001). A series of 4-(2-(substitutedphenyl)-3-phenyl-2,3-dihydro-1H-5-pyrazolyl)-2-methylphenol derivatives and 2-(5-(3-Phenoxyphenyl)-4,5-dihydro (benzoyl)-pyrazol-3-yl) pyridine (**51**) also exhibited good anti-TB activity (Kini et al., 2009).

The search for new anti-TB drugs, compound PA-824 (**52**) as promising anti-TB agents, which has novel mode of actions and efficacy against resistant *Mtb* (Khan et al., 2008). The two diastereomers of 7-methyl-nitroimidazo-oxazine, 7-(*S*)-methyl derivative (**53a, cis**) and the 7-(*R*)-methyl derivative (**53b, trans**) displayed similar activities against *Mtb* with MIC=0.2-0.4 μM (Li et al., 2008). The PA-824 (**52**) along with Metronidazole (**54a**) showed anti-TB effect. Compound **54b** showed promising aerobic inhibition of 99% at 4-8 μM and anaerobic inhibition of 90% at 31.25 μM , which are less than that of PA-824 (Kim et al., 2009). These molecules have shown more potency of aerobic & anaerobic inhibition than parent molecule PA-824. Among all, compound **54b** exhibited aerobic 99% inhibition at 0.039 μM and compound **54c** showed 90% anaerobic inhibition at 1.56-3.13 μM (Kim et al., 2009). Analogues of 2-nitroimidazooxazines, compound **55** has shown best activity with a MIC of 0.11 and 2.7 $\mu\text{g}/\text{mL}$ against *Mtb* in aerobic and anaerobic conditions respectively (Thompson et al., 2009). A imidazo (1,2-*c*) pyrimidine derivatives, compound (**56**) has shown promising MIC of 2 $\mu\text{g}/\text{mL}$ against *Mtb* H37Rv, which is equal to amikacin (Chhabria et al., 2009). The hydrazide derivatives of imidazo (1, 2-*a*) pyrazine, compound **57** showed moderate activity with 86% inhibition at 6.25 $\mu\text{g}/\text{mL}$ (Ozdemir et al., 2009).


A series of 4-(5-cyclobutylloxazol-2-yl) thiosemicarbazones were exhibited *in-vitro* and *in-vivo* activity against *Mtb* H37Rv and MDR-TB (Sriram et al., 2007). The oxazolidinones linezolid (**58**) is a synthetic antibacterial agent, which are active against variety Gram-positive organisms. These compounds inhibit translation at the initiation phase of protein synthesis in bacteria. The compound PNU-100480 and thiomorpholine analogue of linezolid (**59**) showed an interesting anti-TB activity. With this interest, anti-TB activity of oxazolidinones, 3-(1H-pyrrol-1-yl)-2-oxazolidinone analogues of PNU-100480 showed anti-TB activity. A series of nitrofuranyl isoxazolines with increased proteolytic stability over nitrofuranyl amides showed anti-TB activity against *Mtb*. Compound **60a** showed *in-vitro* high potency of 0.00005 $\mu\text{g}/\text{mL}$. However, their *in-vivo* activity was limited by high protein binding and poor distribution. Consequently, a series of non-nitrofuranyl containing isoxazolines had residual anti-TB activity. This led to the discovery of novel isoxazoline **60b** as anti-TB agent, which showed 90% inhibition at a concentration of 1.56 $\mu\text{g}/\text{mL}$ (Tangallapally et al., 2007). A active molecule **60c** has shown a MIC of 0.4 $\mu\text{g}/\text{mL}$ against *Mtb* H37Rv (Rakesh, et al., 2009).




33




34: Ar=4-chlorophenyl




35




36: R=3-phenylpropyl




37: R=4-chlorophenyl





38




39



40

41: R₁=H, R₂=4-F42: R₁=H, R₂=H43: R₁=2-F, R₂=2-F44: R₁=i-C₃H₇, R₂=4-F45: R₁=4-CH₃, R₂=4-F

46



47

The 5-((*E*)-2- arylethenyl)-3-isoxazolecarboxylic acid alkyl ester derivatives were found as promising anti-TB agents. Among all, 5-((*E*)-2-(3,5-Dichloro-4-pyridinyl)ethenyl)-3-

isoxazolecarboxylic acid ethyl ester (**61a**) has shown preminent activity with a MIC 0.59 μ M in MABA assay, whereas compound 5-((*E*)-2-(6-methoxy-4-quinolinyl)ethenyl)-3-

isoxazolecarboxylic acid butyl ester (**61b**) showed the activity against *Mtb* H37Rv with a MIC 1.8 μ M. Both compound showed almost equal potency with standard drugs INH, RMP in terms of activity and cytotoxicity (Pieroni et al., 2009). While in another series, (*R*)-methyl 2-(5-((2-methylbenzo(d)thiazol-5-yl)oxy)methyl)isoxazole-3-carboxamido)-2-phenylacetate (**62**) has shown less activity with a MIC 1.4 μ M in MABA assay (Huang et al., 2009) in comparison to compound **61a**. A series of alkyl 1-heteroaryl-1*H*-1, 2, 3-triazole-4-carboxylates were exhibited anti-TB activity against *Mtb* H37Rv. Among all, the best potency was shown by *n*-pentyl 1-(6-phenylpyridazin-3-yl)-1*H*-1,2,3-triazole-4-carboxylate (**63**) with a MIC of 3.13 μ g/mL (Japelj et al., 2005).

The discovery of anti-TB drug INH of pyridine skeleton prompted the research on piperidine (hexahydropyridine). In this perception, a series of 1-piperidino-3-arylthioureas were evaluated for their anti-TB activity. Compound **64** has shown best potency of MIC 8 μ g/mL against the strain 303 (Hearn et al., 2005). A series of 2-Substituted derivatives of diphenylpyraline and their 1-phenyl and 1-phenethyl analogues were evaluated against *Mtb* H37Rv as well as their cytotoxicity against human cells (HEK-293). Among all, compound **65** showed an inhibition of 75% at a concentration of 6.25 μ g/mL against *Mtb* H37Rv and also found to be least toxic of the series (Weis et al., 2008).

The 2-substituted derivatives of diphenylpyralines (**65**), bamipine (**66a**) and their 1-phenyl analogues exhibited anti-TB activities. Of these, compound **66b** showed best potency of MIC 6.25 μ g/mL (Weis et al., 2008). Where as, piperidinol analogs were exhibited anti-TB activity against *Mtb* H37Rv. Among all, the good potency was shown by compound **67** with a MIC 1.4 μ g/mL and therapeutic index of 13.3 (Sun et al., 2009). The dipiperidine derivatives were discovered as novel anti-TB agents with a hit molecule **68**, which showed preeminent activity with a MIC 7.8, 15.65 μ M in the broth microdilution and BACTEC assay respectively. This compound is also found to less toxic of all with an IC₅₀ 162 μ M against HepG2 cells (Bogatcheva et al., 2010).

A series of 2,6-diarylpiperidin-4-ones and tetrahydropyridin-4-ol based benzimidazole and *o*-arylsulfonyl derivatives exhibited anti-TB activity. Among all, three compounds (**69a**, **69b** and **69c**) have shown equal potency of MIC 16 μ g/mL against *Mtb* H37Rv, which are one-fold more potent than of the standard RIF drug (Aridoss et al., 2008). In 3,5-bis(benzylidene)-4-piperidone (**70**) was found active with a MIC of 0.2 μ g/mL and also found non toxic in mice (Das et al., 2008). In an effort to increase the potency of piperidones, a series of spiro-piperidin-4-ones were evaluated for their anti-TB activity. Among all, compound **71** showed promising *in-vitro* potency of MIC 0.07 μ g/mL and 0.16 μ g/mL against *Mtb* and MDR-TB respectively. Compound **71** also showed *in-vivo* potency by decreasing the bacterial load in lung and spleen tissues with 1.30

and 3.73-log 10 protections respectively, which is comparable to INH (Kumar et al., 2008). A series of *N*-Alkyl-1,2-dihydro-2-thioxo-3-pyridinecarbothioamides were exhibited anti-TB activity against *Mtb* and MAC strains. Among all, compound **72** showed good potency of MIC 0.5 μ g/mL against *Mtb* H37Rv and 2-4 μ g/mL against MAC strains (Pagani et al., 2000). While, the series of 2-pyridine carboxamidrazones (**73**) were showed MIC₅₀ in the range of 160-16 μ g/L against *M. avium* strains (Banfi et al., 2001).

The pyridine derivatives designed as lipophilic precursors were found more active than the unmodified polar isosteres of pyrazinoic acid and nicotinic acid which may be due to better penetration of the compound into the cell wall of *Mtb*. In this view, a series of 1,4-dihydropyridine-3,5-dicarbonyl derivatives with lipophilic groups (**74**) were showed activity against *Mtb* H37Rv, most compound showed 90% inhibition at 2.5 μ g/mL (Desai et al., 2001). A series of pyridines substituted with 1,2,4-oxadiazole-5-ones, 1,2,4-oxadiazole-5-thiones and 1,3,4-oxathiazoline-2-ones exhibited activity against *Mtb* H37Rv. Among all, 1,3,4-oxathiazoline-2-one derivative (**75**) showed best activity with MIC of 4.5 μ g/mL (Gezginci et al., 2001).

A series of heterocyclic chalcones, compound **76** has shown MIC of 6.8 μ g/mL against *Mtb* H37Rv (Lin et al., 2002). A series of substituted *N*-pyridinylsalicylamides (**77**) were exhibited *in-vitro* anti-TB activity against *M. avium* and two strains of *M. kansasii* (Waisser et al., 2004). In the same direction, a series of isonicotinylhydrazones and found a molecule (**78**) active against *Mtb* H37Rv with a MIC of 0.56 μ M, which is more potent than INH (MIC of 2.04 μ M) (Sriram et al., 2005). Another series and found a new derivative (**79**) more active than the former molecule (**78**). It showed equal MIC of 0.49 μ M against *Mtb* H37Rv and INH-resistant *Mtb* strains (Sriram et al., 2006). A series of trans-cinnamic acid derivatives of isonicotinic acid (**80**), showed moderate *in vitro* activity of MIC 3.12 μ g/mL against *Mtb* strain (Carvalho et al., 2008). Whereas, compounds **81a** and **81b** of the isonicotinic acid derivatives developed and showed equal potency of MIC 0.39 μ g/mL against *Mtb* H37Rv and have selectivity index (SI) of >160, which is comparable to INH and better than ciprofloxacin (Imramovsky et al., 2007). A number of (*E*)-*N'*-(monosubstitutedbenzylidene) isonicotinohydrazide derivatives exhibited *in-vitro* anti-TB activity against *Mtb* H37Rv. Five compounds (**82a-e**) have shown significant MIC in the range of 0.31-0.62 μ g/mL in comparison with INH and RIF (Lourenco et al., 2008). Whereas, hybrid of isonicotinic hydrazone of pyrrole (**83**) showed best potency of MIC \leq 0.1 μ g/mL against *Mtb* H37Rv and also has good selectivity index (Bijev. 2006).

Compounds (**84a-f**) showed *in vitro* anti-TB activity in the range of MIC 25-50 μ g/mL (Kumar et al., 2002). In continuation, a number of compounds (**85**) have shown anti-TB potency with a MIC in the range of 12.5-25 μ g/mL (Agarwal et al., 2005a). To

further increase the activity, trisubstituted compound (**86**) (Agarwal et al., 2005b), where the activity was same as compound **85**. Where as, chloro derivatives were found to be highly active against *Mtb*. Compounds (**87a-d**) were found to be active at a MIC of 0.78µg/mL (Agarwal et al., 2002). While, in a series of anilino pyrimidines against *Mtb* H37Ra, the most potent activity was shown by the compound **88** having a MIC of 3.12µg/mL (Morgan et al., 2003).

A series of *N*-phenyl-6-methyl-2-oxo-4-phenyl-1,2,3,4-tetrahydropyrimidine-5- carboxamides were shown anti-TB activity against *Mtb* H37Rv. Two compounds with 2,3-dimethylphenyl (**89a**) and 3,4-dimethyl (**89b**) carbamoyl side chain, respectively, showed 65% and 63% inhibition (Virsoadia et al., 2008). 5-phenyl-6-((3R,4S)-3,4,5-trihydroxypentyl)pyrimidine-2,4-diamine (**90**) selectively inhibited *Mtb* DHFR (El-Hamamsy et al., 2007). Whereas, in a series of methylene-bis-pyrimidinones and methylene-bis-mercapto pyrimidines, pyrimidinone derivative (**91**) shown best potency of 0.1µg/mL while mercapto pyrimidines showed moderate activity (Nagaraj et al., 2007).

Several derivatives of pyrazine nucleus have shown activity against *Mtb*. Pyrazine derivatives substituted with 1,2,4-oxadiazole-5-ones, 1,2,4-oxadiazole-5-thiones and 1,3,4-oxathiazoline-2-ones (Gezginci et al., 2001). The most active compound of the series **92** exhibited a MIC of 4.5µg/mL in comparison to 49µg/mL for pyrazinamide. A series of ring substituted (E)-3-Phenyl-1-(2 pyrazinyl)-2-propen-1-ones were shown efficacy against *Mtb* H37Rv. Among all, compound **93** showed an inhibition of 94% at 12.5µg/mL (Opletalova et al., 2002). While, in a series of pyrazine derivatives, two compound (**94a** and **94b**) have shown equal potency of MIC 6.25 µg/mL against *Mtb* H37Rv (Seitz et al., 2002). Compound **94b** also showed MIC of <0.25 µg/mL against *Mtb* H37Ra.












On the basis of ethionamide, a series of 5-Alkyl-6-(alkyl/aryl sulfanyl) pyrazine-2-carbothioamide, compounds inhibited the growth in the range of 61-91%. Compound (**95**) showed 91% inhibition at a MIC <6.25 µg/mL (Krinkova et al., 2002). While, S-methyl-2-(amino(6-chloropyrazin-2-yl)methylene)hydrazinecarbodithioate (**96**) exhibited moderate potency of MIC 32 µg/mL among simple pyrazine hybrids, against *Mtb* sensitive and wild strains (Foks et al., 2004).

A series of unsubstituted, halogenated and/or alkylated pyrazine-2-carboxylic acid amides connected *via* -CONH- bridge with substituted anilines shown activity against *Mtb* H37Rv. Among all, 5-tert-Butyl-6-chloro-N-(3-trifluoromethylphenyl)pyrazine-2-carboxamide (**97**) has shown the highest activity of MIC 3.13 µg/mL (Dolezal et al., 2008). Compound **98a** and **98b** showed MIC of 0.78, 0.1 µg/mL respectively, against *Mtb* H37Rv. Compound **98a** also showed good activity against atypical strains of *Mtb*(127). In a different approach, A series of 1,4-substituted










piperazine/homopiperazines. Compound **99** showed MIC of 62.5 µM, homopiperazine derivatives (**100a** and **100b**) showed preeminent MIC of 1.56 µM against *Mtb* H37Rv. Compounds **100a** and **100b** also exhibited good selectivity index of 84.6 and 46 respectively (He et al., 2007; Bogatcheva et al., 2006). Homopiperazine compounds (**101a** and **101b**) with a more promising activity of MIC 0.78µg/mL against *Mtb* H37Ra (Zhang et al., 2009). In a different approach pentacycloundecane (PCU) tetra-amine compounds were shown *in-vitro* anti-TB activity against H37Rv and XDR strains of *Mtb* 194. The most active compound (**102**) of the series has shown MIC of 5.04 µM against *Mtb* H37Rv and 1.26 µM against XDR-TB (Onajole et al., 2009).

A series of carbamate derivatives of 1,2-oxazine were shown *in-vitro* anti-TB activity against *Mtb* and *M. lufu* species. Among all compounds, the maximum anti-TB activity was observed for 4,5-dimethyl-2-(*p*-methoxycarbonylamino)phenyl-3,6-dihydro-1,2-oxazine (**103**) (Velikorodov et al., 2006). In a series of 5,6 dihydro-4H-1,3 thiazine derivatives were shown activity against *Mtb* H37Rv, 5-hydroxy-3-phenyl-4-aza-2-thiabicyclo(3.3.1)non-3-ene (**104**) showed 97% inhibition at a concentration of 6.25µg/mL (Koketsu et al., 2002). A series of α -methylene- γ -butyrolactones based on the natural product protolichesterinic acid (**105**) were shown potency against *M. bovis* BCG. Compound **105** was showed improved activity with MICs in the range of 6.25-12.5 µg/mL (Hughes et al., 2005). Several structural analogues of the polyketide passifloricin lactone (**106**) were shown activity against *Mtb* H37Rv (Cardona et al., 2006). Of these, compound **106** exhibited an inhibition percentage higher than 97% at 128µg/mL, while passifloricin A reached 82.9%. Additionally, it has shown best MIC of 17.31µg/mL, which is better than passifloricin A (29.4µg/mL). Peptide deformylase (PDF) is a key enzyme, that deformylates the *N*-formylmethionine, polypeptides, a key step in protein maturation. It is a validating target as an anti-TB agent. In this view, a series of LBK-611 (**107**) derivatives were showed anti-TB activity (Faugeroux et al., 2007).

The purine analogues possessing anti-TB activity have been pursued with great interest. In this perception, 9-benzylpurines with a variety of substituents at 2, 6 or 8 positions were found as good anti-TB agents. High activity was exhibited by 9-benzylpurines carrying a phenyl ethynyl, transstyryl or aryl substituents at the 6th position and generally chlorine at the 2nd position. The most active compounds **108a** and **108b** showed a MIC of 3.13 and 0.78 µg/mL respectively, against *Mtb* H37Rv and also a selectivity index (SI) of 2.7 and 10.4 (Bakkestuen et al., 2000). In continuation, a series of 6-arylpurines having a variety of substituents in the 9 position were screened against *Mtb* H37Rv. The most active compound of the series was again found to be same 9-benzyl-2-chloro-6-(2-furyl)purine (**108b**) having a MIC of 0.78 µg/mL. This compound exhibited



50a: R=H 50b: R=2-Cl
 50c: R=3-Cl 50d: R=4-Cl
 50e: R=2-CH₃



relatively low cytotoxicity and it was also active against several singly drug-resistant strains of *Mtb* (Gundersen et al., 2002). Eleven analogues of 9- sulphonated/sulphenylated 6-mercaptapurines (Scozzafava et al., 2001) and out of them six

exhibited MIC in the range of 0.39-0.78 µg/mL. The most potent compound (**109**) (MIC=0.39 µg/mL) also exhibited good activity against MDR strains of *Mtb*.

Inspired by the above results, a series of 9-aryl-, 9-arylsulfonyl- and 9-benzyl-6-(2-furyl)purines and screened for their anti-TB activity against *M. tuberculosis* H37Rv. Among all, 2-chloro-6-(2-furyl)-9-(4-methoxyphenylmethyl)-9H-purine (**110**) exhibited best potency of MIC 0.39 µg/mL and also low toxicity against mammalian cells and activity inside macrophages (Bakkestuen et al., 2005). In a purine derivatives, 9-(ethylcarboxymethyl)-6-(dodecylthio)-9H-purine (**111**) showed MIC of 0.78 µg/mL (Pathak et al., 2004). In the analogues of agelasine E (**112a**), one derivative (**112b**) showed promising activity with MIC of 1.56 µg/mL against *Mtb* H37Rv (Bakkestuen et al., 2005). After the keen observation of SAR of above molecules, 6-(2-furyl)-9-(p-methoxybenzyl)purines carrying a variety of substituents in the 2- or 8-position and was successful in identifying a more potent molecule (**113**, MIC=0.20 µg/mL) (Braendvang et al., 2007) of above all series. The purine derivatives, found a more potent molecule (**1114**) of above all series, which has shown an IC90 of <0.20 µg/mL against *Mtb* H37Rv (Braendvang et al., 2009).


In search of new anti-TB purine type analogues, a series of 1-(1-(4-hydroxybutyl)-1,2,3-triazol-4 and 5)-ylmethyl)-1H pyrazolo(3,4-d)pyrimidines and all of them were inactive but one compound (**115**) has shown MIC of 12.5 µg/mL (Moukha-Chafiq et al., 2001; Moukha-Chafiq et al., 2002; Moukha-Chafiq et al., 2007).

In continuation, a series of di/trisubstituted pyrazolo(3,4-d)pyrimidines (**116**, **117**) and observed no significant anti-TB activity at concentrations up to 6.25 µg/mL. A series of *N,S*-bis-alkylated thiopyrazolo(3,4-d)pyrimidines, based on sequential *S*-then *N*-alkylation, is carried out. These compounds showed significant anti-TB activity (MICs down to ≤2 µg/mL) and their potential as significant drug-like leads is substantiated through cytotoxicity evaluation and in silico profiling. Among all, one compound (**118**) has shown MIC=0.5-1 µg/mL against *Mtb* H37Rv (Ballell et al., 2007). A homologous series of three pyrazolopyrimidine analogues (**119a-c**) of a hypothetical intermediate in the lumazine synthase-catalyzed reaction and evaluated as lumazine synthase inhibitors. All three compounds were extremely potent inhibitors (Inhibition constant: K_i =15-40 nM) of the lumazine synthases of *Mtb* with inhibition constants in the low nanomolar to subnanomolar range. Molecular modeling of one of the homologues bound to *Mtb* lumazine synthase suggests that both the hypothetical intermediate in the lumazine synthase-catalyzed reaction pathway and the metabolically stable analogues bind similarly (Zhang et al., 2007). In a series of Thieno(2,3-d)pyrimidin-4-one, two compounds (**120a** and **120b**) have shown moderate potency of 5µM/L against *Mtb* and *M. avium*, which is equal to that of rifampicin (Chambhare et al., 2003).


The identification of new promising quinoline based anti-TB agents, 2,8-dicyclopentyl-4-methylquinoline (DCMQ, **121**)

(Jain et al., 2003) and Diarylquinoline (TMC207, **122**) (181) have definitely initiated the optimization of quinoline for anti-TB drugs. In this concern, 1-(5-isoquinolinesulfonyl)-2-methylpiperazine (**123**), a protein kinase inhibitor for its anti-TB profile and found to inhibit the growth of two different mycobacterial strains, the slow-growing *M. bovis* Bacille Calmette Guerin (BCG) and the fast-growing saprophyte *M. smegmatis* mc2 155, in a dose-dependent manner. While screening for the effect of kinase inhibitors on mycobacterial growth, millimolar concentrations of **123** induced a 40% decrease in the growth of *M. bovis* BCG when measured as a function of oxidative phosphorylation. This **123**-induced decrease in growth was shown to involve a 2-log fold decrease in the viable counts of *M. smegmatis* within a 48h period and a 50% reduction in the number of BCG viable counts within a 10-day period. Micromolar concentrations of **123** induced a significant decrease in the activity of the *M. tuberculosis* protein serine/threonine kinase (PSTK) PknB. The inhibition of mycobacterial growth as well as the inhibition of a representative *Mtb* protein serine/threonine kinase PknB suggests that conventional PSTK inhibitors can be used to study the role that the mycobacterial PSTK family plays in controlling bacterial growth (Gaurrand et al., 2006; Drews et al., 2001). A series of quinolinyl hydrazones and majority of the tested compounds showed an inhibitory activity between 95 and 100%. The most potent compounds of the series (**124a-c**) were having a MIC of 0.78µg/mL. These results indicated that the activity was significantly affected by substituents both on the quinoline nucleus and hydrazinoic moiety. On quinoline nucleus the most effective substituents resulted were 6-cyclohexyl, 7-methoxy or ethoxy and 7-chloro groups. Similarly, for the hydrazinoic moiety greater effectiveness resulted for *para* and ortho-methoxynaphthyl substituents whereas disubstitution with chlorine resulted in inactive compounds (Savini et al., 2002). Inspired with the activity profile of DCMQ (**121**), four new series of the ring-substituted quinolinecarbohydrazides were evaluated. Of these 3-quinolinehydrazides and *N2*-alkyl/*N2,N2*-dialkyl/*N2*-aryl-4-(1-adamantyl)-2-quinolinecarboxamides showed moderate activity of MIC in the range of 6.25-3.125 µg/mL against *Mtb* H37Rv. The most active compounds were adamantyl derivatives (**125a** and **184b**) exhibited MIC of 3.125 µg/mL (Monga et al., 2004). Whereas, a number of thirty-three quinoline derivatives based on TMC207 (**122**) and found a molecule (**126**) active with a MIC=3.12 µg/mL (de Souza et al., 2009). With the same interest, 3-benzyl-6-bromo-2-methoxy-quinolines and amides of 2-((6-bromo-2-methoxy-quinolin-3-yl)-phenylmethyl)-malonic acid monomethyl ester. Among all, four compounds (**127a-d**) showed moderate activity of MIC 6.25 µg/mL against *Mtb* H37Rv (Upadhayaya et al., 2009).


Recently, a series of substituted quinolinyl chalcones and substituted quinolinyl pyrimidines were evaluated for their in




58




59




60a: R=

60b: R= COOC₂H₅


60c




61a




61b




62




63




64




65




66a




66b




67



68



















69a



69b


in vitro anti-TB activity against *Mtb* H37Rv. Chalcone derivatives **128a** and **128b** have shown antitubercular activity of MIC 3.12 $\mu\text{g/mL}$ and were nontoxic against VERO, MBMDM cell lines (Sharma et al., 2009). The anti-TB potential of NAS-91 (**129**) and found that NAS-91 has multiple targets, which is particularly

desirable for avoiding the emergence of resistant strains of *Mtb*. Therefore, NAS-91 represents a potent pharmacophore and appears to be a promising lead compound for future inhibitor development against TB (Gratraud et al., 2008). The quinoline-3-carbohydrazone derivatives were screened for their anti-TB



efficacy. Among all, two compounds (**130a** and **130b**) have shown promising activity with a MIC 0.625, 2.5 and 1.25 $\mu\text{g}/\text{mL}$ against *Mtb* H37Rv, *M. smegmatis* and *M. fortuitum* respectively. These compounds have shown almost equal potency similar to

that of standard rifampicin (Eswaran et al., 2010). Whereas in the series of 4-quinolyldrazones, the most active compound (**131**) displayed an anti-TB activity of MIC 0.6 μM and selectivity index 2.27 (Gemma et al., 2009).



84 a-f,

85 and 86,

a: R=*N*-phenylpiperazine, R'=4-fluorobenzene


b: R=*c*-heptylamine, R'=4-methylbenzene

c: R=Octylamine, R'=3-pyridine

d: R=*n*-propylamine, R'=3-pyridine

e: R=4-(2- aminoethyl)morpholine, R'=3-pyridine

f: R=Octylamine, 130a-f R'=4-methoxybenzene




87a-d

a: R=*n*-C₃H₇, Ar=2-Theinyl


b: R=4-Cl C₆H₄CH₂, Ar=2-Theinyl

c: R=C₃H₇, Ar=3-F C₆H₄


d: R=*n*-C₃H₇, Ar=2-Furyl




88




89a




89b




90




91



92




93




94a: R=4-nitrobenzyloxy


94b: R=4-acetoxybenzyloxy



95



96



97

A similar approach, 7-chloro-4-quinolinylhydrazone derivatives and found three molecules (**132a-c**) with a moderate anti-TB activity with a MIC 2.5µg/mL. These compounds were found to be nontoxic against J774 cell line up to the

concentration 100 µg/mL (Candéa et al., 2009). Quinoline derivatives consisting of triazolo, ureido and thioureido substituents at C-6 position, ureido derivative (**133a**) and triazolo derivative (**133b**) have shown moderate activity of MIC 3.125

$\mu\text{g/mL}$ against *Mtb* H37Rv (Upadhayaya et al., 2009). A series of amino acid conjugates of 4-(adamantan-1-yl) group containing quinolines. The most active nontoxic compound (**134**) of the series exhibited increased potency of $1\mu\text{g/mL}$ against *Mtb* H37Rv and $3.125\mu\text{g/mL}$ against MDR strain, in comparison to former molecules (**133a** & **133b**) (Nayyar et al., 2009). In the same direction of approach, quinoline-based derivatives were evaluated for their anti-TB efficiency. Among all, compound **135** has shown remarkable activity of MIC $0.77\mu\text{M}$ against *Mtb* H37Rv and $0.99\text{--}1.55\mu\text{M}$ against drug-resistant strains (Lilienkampf et al., 2009). In continuation the same group synthesized isoxazole based quinoline derivatives and found a lead molecule **136**, which showed MIC of $0.2\mu\text{M}$ and $2.6\mu\text{M}$ in MABA and LORA assay against *Mtb* H37Rv (Mao et al., 2009). Thus, optimization of quinolines for the development of anti-TB agents is a fruitful approach.

The quinoxaline derivatives show very interesting antimicrobial properties and recently, some researchers have identified the anti-TB activities of various 2-methylquinoxaline 1,4-dioxides, confirming that the presence of a methyl (or halogenomethyl) group at 2(3) position of this ring (**137a** and **137b**) is favourable for antimicrobial activity. In this context as contribution in the development of quinoxaline derivatives, a number of 36, 6(7)-substituted-3-methyl- or 3-halogenomethyl-2-phenylthio-phenylsulphonyl-chloro-quinoxaline 1,4-dioxides were screened for their in vitro anti-TB activity. Among all, two compounds **138a** and **138b** exhibited great potency of MIC $0.39\mu\text{g/mL}$ against *Mtb*, which is comparable to Rifampicin (MIC= $0.25\mu\text{g/mL}$) (Carta et al., 2002). In another series, four compounds (**139a-d**) have shown moderate activity of MIC $2\mu\text{g/mL}$ (Carta et al., 2004). In the series of quinoxaline derivatives, lack of 1,4-dioxide showed reduction in the activity. Most active compound (**140**) showed MIC of $6.25\mu\text{g/mL}$ against MTB H37Rv and $0.5\mu\text{g/mL}$ against MTB H37Ra (Seitz et al., 2002). Which prompted to continue the optimization of quinoxaline 1,4-dioxide.

Inspired with the activity profile of **137a** and **137b**, (Zarranz et al., 2003) a series of quinoxaline-2-carboxamide 1,4-di-*N*-oxide derivatives and evaluated for their in vitro anti-TB activity against *Mtb* H37Rv. Among all, compound **141a** exhibited best MIC of $0.78\mu\text{M}$, has a solubility problem, while compound (**141b**) having MIC of $3.13\mu\text{M}$ has a best selectivity index (SI= >40.06). A series of quinoxaline 1,4-di-*N*-oxide derivatives by varying the 2-position and found that 2-methylquinoxaline 1,4-di-*N*-oxides (**142a** and **142b**) were most active of the series with a MIC of $0.39, 0.78\mu\text{M}$ respectively and also have better selectivity index (8.46, 20.43) (Jaso et al., 2003). Villar et al. also found that the compound **142b** is also active against resistant strains of *Mtb* (Villar et al., 2008). In another series, 2-benzyl-3-(methoxycarbonyl)quinoxaline 1,4-dioxide (**143**) has shown best

potency above all, with a MIC= $0.10\mu\text{g/mL}$ and selectivity index SI= 470 (Jaso et al., 2005).


A new series of 3-phenylquinoxaline 1,4-di-*N*-oxide having selectivity against *Mtb* have been evaluated. 34 out of the 70 tested compounds showed an MIC value less than $0.2\mu\text{g/mL}$, a value on the order of the MIC of rifampicin. Furthermore, 45% of the evaluated derivatives showed a good in vitro activity/toxicity ratio. The most active compound was 7-methyl-3-(4'-fluoro)phenylquinoxaline-2-carbonitrile 1,4-di-*N*-oxide (**144**) (MIC $<0.2\mu\text{g/mL}$ and SI >500) (Vicente et al., 2009). In conclusion, the potency, low cytotoxicity and selectivity of these compounds make them valid lead compounds for synthesizing new anti-TB agents.

Benzothiadiazine and pyranopyridine derivatives

In an effort to develop new and more effective therapies to treat tuberculosis, Kamal et al. synthesized a series of benzothiadiazine 1,1-dioxide derivatives and their in vitro activity was evaluated against *Mtb*, *M. avium* and *M. intracellulare*. Of these, compound **145** showed best potency of MIC $0.5\mu\text{g/mL}$ against MTB H37Rv and $0.5\text{--}2\mu\text{g/mL}$ against resistant strains. However, the in vivo testing in a mouse model of TB infection did not show significant anti-TB activity, probably because of its poor bioavailability (Kamal et al., 2006). In continuation, 5-nitrofuran, 5-nitrothiophene and arylfuran coupled benzothiadiazines and on evaluation, these compounds exhibited moderate anti-TB activity. The most active compound (**146**) displayed a MIC of $1\mu\text{g/mL}$ against MTB H37Rv (Kamal et al., 2007). With the same inspiration, a number of fifteen 2-amino-6-methyl-4-aryl-8-((E)-arylmethylidene)-5,6,7,8-tetrahydro-4*H*-pyrano(3,2-*c*)pyridine-3-carbonitriles and evaluated for their anti-TB activity. Among all, compound **147** was found to be the most potent compound (MIC: $0.43\mu\text{M}$) against MTB and MDR-TB, being 100 times more active than isoniazid against MDR-TB (Kumar et al., 2007).

A Series of 9-substituted tetrahydroacridines for their anti-TB activity. These derivatives exhibited promising activity profile, with MIC in the range of $6.25\text{--}0.78\mu\text{g/mL}$ against *Mtb*. Compound **148a** displayed best MIC of $0.78\mu\text{g/mL}$ against MTB-H37Rv, while compound **148b** displayed preeminent activity against MTB-H37Ra with MIC $1.56\mu\text{g/mL}$ (Tripathi et al., 2006).

On this basis and to minimize the side-effects and to improve the anti-TB activity of Clofazimine (**149**), 3-(2,4-dichloroanilino)-10-(2,4-dichlorophenyl)-2,10-dihydro-2-(2,2,6,6-tetramethyl piperid-4-ylimino)phenazine (B4128) (**150**), which posses a similar mode of action of Clofazimine (Reddy et al., 1999; Matlola et al., 2001). With the same motivation, a series of phthalimido- and naphthalimido-linked phenazines and found two compounds (**151a** and **151b**) with a potency of MIC 1



$\mu\text{g/mL}$ against *M. tuberculosis* H37Rv. These compounds also exhibited potency against resistant strains of *Mtb* (Kamal et al., 2005). Whereas in a series of phenazine carboxamides, compounds **152a** and **152b** showed excellent activity against *Mtb* H37Rv with a MIC of 0.19 $\mu\text{g/L}$ and also against drug-resistant strains of *Mtb*. Most interestingly, this series was found to be

nontoxic (De Logu et al., 2009), validating them as future anti-TB drug candidates.

Phenothiazines have been reported for their anti-TB activity for many years, and the phenothiazine drug chlorpromazine (CPZ) (**153a**) is reported to have been successfully used to treat a TB patient. In this concern, a series of psychotropic

phenothiazines were examined as anti-TB agents against *Mtb* H37Rv. Among all, three compounds (**153b-d**) exhibited promising activity with a mean MIC of 2.13 µg/mL (Madrid et al., 2007). Whereas quaternized CPZ, triflupromazine (**154a**) and promethazine (**154b**) derivatives inhibited non-replicating *Mtb* at concentrations equal to or double their MICs against the actively growing strain. All the active compounds (**154c-f**) were non-toxic toward Vero cells (IC₅₀ > 128 µM). Based on SAR it was concluded that the benzyl or substituted benzyl groups, an electron-withdrawing substituent on the phenothiazine ring improved the potency. Commonly the optimum anti-TB structures possessed *N*-(4- or 3-chlorobenzyl) substitution on triflupromazine (Bate et al., 2007). While a macrolactone (**155**) derived from benzo(a)phenazine exhibited best potency against *Mtb* H37Rv with a MIC 0.62 µg/mL, which is better than that of Rifampacin (Silva et al., 2009).

In search of potential anti-TB agents, pyridazinoindole analogues screened for inhibition of the growth of *Mtb*. The most active compound (**156**) exhibited a MIC₅₀ of 1.42 µg/mL against *Mtb* H37Rv (MTB) (Velezheva et al., 2004). In the series (2-aryl-3,4-dihydro-2H-thieno(3,2-b)indoles), compound **157** was found to be the most active compound with MIC of 0.4 µg/mL against MTB and MDR-TB (Karthikeyan et al., 2009).

With the same motivation, a series of pyrrolo(1,2-a) quinoxaline-2- or -4-carboxylic acid hydrazides and one compound (**158**) showed an interesting activity at 6.25 µg/mL against *Mtb* H37Rv, with a 100 percentage inhibition (Guillon et al., 2004). Compound **159** inhibited 80% at a concentration of 6.25 µM (Scozzafava et al., 2001). While, Enamine-containing analogues of heteroarylquinones showed promising activity with a MIC in the range 6.25-0.1 µg/mL against *Mtb* H37Rv. The best selectivity index (SI=15.1) was displayed by the molecule (**160**) with a MIC 0.39 µg/mL (Copp et al., 2005).

The structure-based design in the discovery of alkyl substituted diphenyl ether, inhibitors of InhA, the enoyl reductase from *Mtb*. However, despite their promising in vitro activity, these compounds have Clog_P values of over 5. In efforts to reduce the lipophilicity of the compounds, and potentially enhance compound bioavailability, A series of substituted hetero/aryl ethers and of these, one compound (**161**) exhibited a moderate MIC₉₀ 3.13 µg/mL but have improved Clog_P value (am Ende et al., 2008). A series of pthalamide derivatives, compound (**162**) displayed a MIC of 5 µg/mL against *Mtb* H37Rv and a good selectivity index (Jean et al., 2009). Series of thiophene (**163**) and benzopyrrole/pyridine (**164a** and **164b**) triarylmethanes (Parai et al., 2008; Panda et al., 2007), thiophene analogues displayed MIC in the range 3.12-12.5 µg/mL and benzopyrrole/pyridine analogues displayed 6.25-25 µg/mL against *Mtb* H37Rv.




















A series of spiro-pyrrolothiazoles, the best potency was displayed by compound **165** with a MIC of 0.6 µM against *Mtb*

and MDR-TB (Karthikeyan et al., 2010). Whereas 4-(morpholin-4-yl)-N'-(arylidene) benzohydrazide derivative (**166**) which showed an inhibition of 96.78% at a concentration of 0.05 µg/mL against *Mtb* H37Rv. This compound also showed good percentage of inhibition against clinical isolates of MDR strains (Raparti et al., 2009). The 3-amino-imidazo(1,2-a)pyridines as a novel class of *Mtb* glutamine synthetase inhibitors. The compound (**167**) showed an inhibition of IC₅₀ = 0.38±0.02 µM (Odell et al., 2009)

In search of novel anti-TB agents, tetrahydroindazole based compounds and evaluated their efficiency (Guo et al., 2010). The 1,3-benzothiazin-4-ones (BTZ) kills *Mtb* by blocking arabinan synthesis. The most advanced compound, BTZ043 (**168**), was found to a candidate for inclusion in combination therapies for both drug-sensitive and extensively drug-resistant TB (Makarov et al., 2009; Abdel-Rahman et al., 2009).














DISCUSSION

Despite these positive changes there are still problems that need to be tackled. A critical question today is whether they are sufficient to bring improved treatment to patients in the next few years. A first challenge concerns the sustainability of the current effort. The next important question is whether there are a sufficient number of promising compounds in the TB pipeline for a broadly effective new treatment combination to be developed. Although different attrition rates might apply, the number of candidate compounds is still small compared to the drug pipelines for diseases of major concern to wealthy countries such as cancer or cardiovascular diseases (and the number of companies engaged in the latter is also greater). Furthermore, many of the compounds in the pipeline are either derivatives of existing compounds or they target the same cellular processes as drugs currently in use. Whilst analogues and derivatives are far quicker to develop, they may be subject to cross-resistance, as has been the case with the new rifamycins and quinolones. Modern technologies and rational approaches to drug design (such as creation of genomic libraries of *Mtb* conditional knock-out mutants for comprehensive target identification and validation, target-based drug discovery, or determination of three dimensional crystal structure of molecular targets) are still weakly implemented in the field of drug discovery for TB. Even the more promising candidate compounds currently in clinical development were identified serendipitously in screenings that were not designed originally for activity against *Mtb*. There is consensus among the TB scientific community that in order to obtain a real breakthrough in TB therapy and drastically shorten treatment there is an urgent need for rational approaches aimed at tackling the problem of mycobacterial persistence. The adaptations that allow *Mtb* is to persist in the host despite a vigorous adaptive immune response likely contribute to the



difficulty in curing TB with current chemotherapy. Although drugs currently in the pipeline could significantly shorten treatment, it is likely to remain a matter of months rather than





weeks or days. There are two major roadblocks that hamper the implementation of rational drug design in TB drug discovery. The first is the lack of a comprehensive characterization of the

**127a:** R=Imidazolyl**127b:** R=C-(Thiophen-2-yl)-methylamine**127c:** R=Pyrazolyl**127d:** R=6-Amino-chromen-2-one**128a:** R=2,3-dimethoxy**129****128b:** R=2,5-dimethoxy**130a****130b****131****132 a:**R=Br **b:**R=F **c:**R=OCH₃**133a****133b****134****135****136**


fundamental biology of mycobacteria as they persist in human tissues, which prevents the identification and validation of potential targets that are relevant for the survival of the bacteria in vivo. The second is the weak engagement into early-stage drug discovery; as a consequence the advanced knowledge about M.

tuberculosis metabolism, physiology and genetics is not being translated into validated targets that can be used for screening of new lead compounds (Asif. 2012a; Asif. 2012b).


As part of the Grand Challenges in Global Health initiative the Gates Foundation is funding research into the molecular

137a: R₁=H137b: R₁=Br138a: R₁=H, R₂=Cl138b: R₁=CF₃, R₂=H139a: R₁=H, R₂=F139b: R₁=H, R₂=OCH₃139c: R₁=R₂=OCH₃

139d




140




141a: R=Cl


141b: R=H



142a: R=Cl 142b: R=H



143





















144

pathways of persistence, with the aim of novel target identification. In addition, the Gates Foundation recently announced a new initiative that specifically aims at accelerating drug discovery for tuberculosis. While acknowledging this significant contribution, it is important not to rely exclusively on a single initiative to address a complex scientific problem of such great importance. Much attention must be paid to these critical issues. If faster progress is to be achieved in drug discovery for TB then the advanced knowledge about *Mtb* metabolism and physiology needs to be translated into validated targets that can be used for screening of new lead compounds. A key difficulty lies in securing sustained funding for translational research projects such as target validation and chemical genetics. Rare exceptions are made for occasional grants based on request for application, but generally it is very difficult for academic labs to obtain funds for projects that fall between the areas of basic and applied research. The private sector for its part is reluctant to engage in early stage drug discovery projects; drug development is instead only embarked upon when rigorously validated targets are available or a lead compound has been already identified. Real improvements in TB treatment will require substantial strengthening of early-stage discovery research to identify new compounds and targets (Asif. 2012c; Asif et al., 2011; Asif, et al., 2012; Asif et al., 2013). Without a thriving background of


discovery-oriented translational research, which is largely dependent on public funding, drug development is destined to fail in terms of long-term goals for effective TB management. Existing modern technologies need to be urgently and more comprehensively applied to TB if the pipeline for drug R&D is to be filled. The reluctance of the pharmaceutical sector to invest in early-stage discovery research for neglected diseases exacerbates the pressing need to translate basic scientific knowledge into novel targets and fresh approaches towards improved therapies. Without proper public engagement in early stage drug discovery and implementation of rational approaches, progress in innovation will be severely hindered.

FUTURE PERSPECTIVE

The unremitting and steady rise in TB together with the emergence of resistance against traditional anti-TB drug regimen and the pathogenic synergy with HIV has put enormous pressure on public health systems to introduce new treatments. In MDR-TB, it is important to understand how the resistance emerges. Consequently, great efforts have been made in the area of *Mtb* genomics, proteomics and target identification via advanced technologies and therefore several welcome developments comes in the light having novel target with newer mode of action.



Remarkably, the mechanisms of action of these new drugs are well-understood with new and novel target. Further investment in



developing fundamental genetic systems and more accurate models of human disease would significantly facilitate TB drug discovery efforts in the long term, in particular enabling robust validation of novel targets.

CONCLUSION

Tuberculosis (TB) is a chronic infectious disease caused by *Mtb*. The term MDR-TB is used to describe strains that are resistant to two or more of the five first-line anti-TB drugs. Treatment regimen of TB comprises five first line antiTB drugs namely isoniazid, rifampicin, pyrazinamide, streptomycin and ethambutol followed by second line antiTB drugs namely fluoroquinolones and one of the injectable aminoglycosides. Besides the traditional anti-TB drugs available commercially, several new heterocycles were synthesized in recent past. The new potential anti-TB agents have been classified according to their chemical entities. The new developed and more effective molecules are also effective against MTB and MDR-TB.

REFERENCES

- Abdel Abdel-Rahman, H.M., El-Koussi, N.A., Hassan, H.Y., 2009. Fluorinated 1,2,4-Triazolo(1,5-a)pyrimidine-6-carboxylic Acid Derivatives as Antimycobacterial Agents. *Archiv der Pharmazie-Chemistry in Life Sciences* 342(2), 94-99.
- Agarwal, A., Srivastava, K., Puri, S.K., Sinha, S., Chauhan, P.M.S., 2005. A small library of trisubstituted pyrimidines as antimalarial and antitubercular agents. *Bioorganic & Medicinal Chemistry Letters* 15(23), 5218-5221.
- Agarwal, A., Srivastava, K., Puri, S.K., Sinha, S., Chauhan, P.M.S., 2005. Solid support synthesis of 6-aryl-2-substituted pyrimidin-4-yl phenols as anti-infective agents. *Bioorganic & Medicinal Chemistry Letters* 15(22), 4923-4926.
- Agarwal, N., Srivastava, P., Raghuvanshi, S.K. et al., 2002. Upadhyay Chloropyrimidines as a new class of antimicrobial agents. *Bioorganic & Medicinal Chemistry* 10(4), 869-874.
- Alcaide, F., Calatayud, L., Santín, M., Martín, R., 2004. Comparative In Vitro Activities of Linezolid, Telithromycin, Clarithromycin, Levofloxacin, Moxifloxacin, and Four Conventional Antimycobacterial Drugs against

- Mycobacterium kansasii*. Antimicrobial Agents & Chemotherapy 48(12), 4562-4565.
- am Ende, C.W., Knudson, S.E., Liu, N. et al., 2008. Synthesis and in vitro antimycobacterial activity of B-ring modified diaryl ether InhA inhibitors. Bioorganic & Medicinal Chemistry Letters 18(10), 3029-3033.
- Aridoss, G., Amirthaganesan, S., Kumar, N.A. et al., 2008. A facile synthesis, antibacterial, and antitubercular studies of some piperidin-4-one and tetrahydropyridine derivatives. Bioorganic & Medicinal Chemistry Letters 18(24), 6542-6548.
- Arya, K., Agarwal, M., 2007. Microwave prompted multigram synthesis, structural determination, and photo-antiproliferative activity of fluorinated 4-hydroxyquinolinones. Bioorganic & Medicinal Chemistry Letters 17(1), 86-93.
- Ashraf, M.W., Bilal, M., Iqbal, M., 2015. Comparative analysis of antiglycation potential of vegetables aqueous and methanolic extracts. Current Science Perspectives 1(1), 1-15.
- Asif, M., 2012. A Review of Antimycobacterial Drugs in Development. Mini review in Medicinal Chemistry 12(13), 404-418.
- Asif, M., 2012. Study of clinically used and recently developed antimycobacterial agents. Oriental Pharm Exp Med, 12:15-34.
- Asif, M., 2012. Study of currently used antimycobacterials, their analogues and recently developed agents. Indian Drugs 49(7): 5-19.
- Asif, M., 2015a. Pharmacologically potentials of different substituted coumarin derivatives. Chemistry International 1(1), 1-11.
- Asif, M., 2015b. Chemistry and antioxidant activity of plants containing some phenolic compounds. Chemistry International 1(1), 35-52.
- Asif, M., 2015c. 7Antiviral and antiparasitic activities of various substituted triazole derivatives: A mini review. Chemistry International 1(2), 71-80.
- Asif, M., 2015d. Role of some nutritional complements and biological supplements in the management of epilepsy. Current Science Perspectives 1(1), 1-11.
- Asif, M., 2015e. Antiglycation activity of vegetables aqueous and methanolic extracts. Current Science Perspectives 1(1), 33-40.
- Asif, M., 2015f. The impact of dietary fat and polyunsaturated fatty acids on chronic renal diseases. Current Science Perspectives 1(2), 51-61.
- Asif, M., Siddiqui, A.A., Husain, A., 2012. Quinolone derivatives as antitubercular drugs. Medicinal Chemistry Research, DOI 10.1007/s00044-012-0101-3.
- Asif, M., Singh, A., Lakshmayya., 2013. The development of structurally different new antitubercular molecules containing pyridazine ring system. Chronicle of Young Scientist 4(1), 1-8.
- Asif, M., Singh, A., Ratnakar, L., 2011. Antimicrobial Agents: Brief Study of Pyridazine Derivatives against Some Pathogenic Microorganisms. Journal of Pharmacy Research 4(3), 664-667.
- Bakkestuen, A.K., Gundersen, L.L., Langli, G., Liu, F., Nolsøe, J.M.J., 2000. 9-Benzylpurines with Inhibitory Activity against *Mycobacterium tuberculosis*. Bioorganic & Medicinal Chemistry Letters 10(11), 1207-1210.
- Bakkestuen, A.K., Gundersen, L-L., Petersen, D., Utenova, B.T., Vik, A., 2005. Synthesis and antimycobacterial activity of agelasine E and analogs. Org Biomol Chem 3(6), 1025-1033.
- Bakkestuen, A.K., Gundersen, L-L., Utenova, B.T., 2005. Synthesis, Biological Activity, and SAR of Antimycobacterial 9-Aryl-, 9-Arylsulfonyl-, and 9-Benzyl-6-(2-furyl)purines. Journal of Medicinal Chemistry 48(7), 2710-2723.
- Ballell, L., Field, R.A., Chung, G.A.C., Young, R.J., 2007. New thiopyrazolo(3,4-d)pyrimidine derivatives as antimycobacterial agents. Bioorganic & Medicinal Chemistry Letters 17(6), 1736-1740.
- Banfi, E., Mamolo, M.G., Zampieri, D., Vio, L., Bragadin, C.M., 2001. Antimycobacterial activity of N1-{1-(3-aryl-1-(pyridin-2-,3- or 4-yl)-3-oxo)propyl}-2-pyridine carboxamides. J Antimicrob Chemother. 48(5), 705-707.
- Barry, CE3rd., Slayden, R.A., Sampson, A.E., Lee, R.E., 2000. Use of genomics and combinatorial chemistry in the development of new antimycobacterial drugs. Biochem Pharmacol 59(3), 221-231.
- Bastian, I., Colebunders, R., 1999. Treatment and prevention of multidrug-resistant tuberculosis. Drugs, 58(4), 633-661.
- Bate, A.B., Kalin, J.H., Fooksman, E.M., et al., 2007. Synthesis and antitubercular activity of quaternized promazine and promethazine derivatives. Bioorganic & Medicinal Chemistry Letters 17(5), 1346-1348.
- Biava, M., Porretta, G.C., Deidda, D., Pompei, R., Tafi, A., Manetti, F., 2004. Antimycobacterial compounds. New pyrrole derivatives of BM212. Bioorganic & Medicinal Chemistry 12(6), 1453-1458.
- Biava, M., Porretta, G.C., Deidda, D., Pompei, R., Tafi, A., Manetti, F., 2003. Importance of the Thiomorpholine Introduction in New Pyrrole Derivatives as Antimycobacterial Agents Analogues of BM 212. Bioorganic & Medicinal Chemistry 11(4), 515-520.
- Biava, M., Porretta, G.C., Poce, G., et al., 2005. Antimycobacterial compounds. Optimization of the BM212

- structure, the lead compound for a new pyrrole derivative class. *Bioorganic & Medicinal Chemistry* 13(4), 1221-1230.
- Biava, M., Porretta, G.C., Poce, G., et al., 2006. Antimycobacterial agents. Novel diarylpyrrole derivatives of BM212 endowed with high activity toward *Mycobacterium tuberculosis* and low cytotoxicity. *Journal of Medicinal Chemistry* 49(16), 4946-4952.
- Biava, M., Porretta, G.C., Poce, G., et al., 2008. 1,5-Diphenylpyrrole Derivatives as Antimycobacterial Agents. Probing the Influence on Antimycobacterial Activity of Lipophilic Substituents at the Phenyl Rings. *Journal of Medicinal Chemistry* 51(12), 3644-3648.
- Biava, M., Porretta, G.C., Poce, G., et al., 2009. 1,5-Diaryl-2-ethyl pyrrole derivatives as antimycobacterial agents: Design, synthesis, and microbiological evaluation. *European Journal of Medicinal Chemistry* 44(11), 4734-4738.
- Bijev, A., 2006. New Heterocyclic Hydrazones in the Search for Antitubercular Agents: Synthesis and In Vitro Evaluations. *Letters in Drug Design & Discovery*, 3(7), 506-512.
- Bishai, W.R., Chaisson, R.E., 1997. Short-course chemoprophylaxis for tuberculosis. *Clin. Chest Med.* 18(1), 115-122.
- Bogatcheva, E., Hanrahan, C., Chen, P., et al., 2010. Discovery of dipiperidines as new antitubercular agents. *Bioorganic & Medicinal Chemistry Letters* 20(1), 201-205.
- Bogatcheva, E., Hanrahan, C., Nikonenko, B., et al., 2006. Identification of New Diamine Scaffolds with Activity against *Mycobacterium tuberculosis*. *J. Med. Chem.* 49(11), 3045-3048.
- Braendvang, M., Bakkenc, V., Gundersen, L-L., 2009. Synthesis, structure, and antimycobacterial activity of 6-(1(3H)-isobenzofuranylidene-methyl)purines and analogs. *Bioorganic & Medicinal Chemistry* 17(18), 6512-6516.
- Braendvang, M., Gundersen, L-L., 2007. Synthesis, biological activity, and SAR of antimycobacterial 2- and 8-substituted 6-(2-furyl)-9-(p-methoxybenzyl)purines. *Bioorganic & Medicinal Chemistry* 15(22), 7144-7165.
- Candéa, A.L.P., Ferreira, M.L., Pais, K.C., et al., 2009. Synthesis and antitubercular activity of 7-chloro-4-quinolinylhydrazones derivatives. *Bioorganic & Medicinal Chemistry Letters* 19(22), 6272-6274.
- Cardona, W., Quinones, W., Robledo, S., et al., 2006. Antiparasite and antimycobacterial activity of passifloricin analogues. *Tetrahedron*, 62(17), 4086-4092.
- Carta, A., Loriga, M., Paglietti, G., et al., 2004. Synthesis, antimycobacterial, antitrichomonas and anti-candida in vitro activities of 2-substituted-6,7-difluoro-3-methylquinoxaline 1,4-dioxides. *European Journal of Medicinal Chemistry* 39(2), 195-203.
- Carta, A., Paglietti, G., Nikookar, M.E.R., Sanna, P., Sechi, L., Zanetti, S., 2002. Novel substituted quinoxaline 1,4-dioxides with in vitro antimycobacterial and anticandida activity. *European Journal of Medicinal Chemistry* 37(5), 355-366.
- Carta, A., Palomba, M., Paglietti, G., Molicotti, P., Paglietti, B., Cannas, S., Zanetti, S., 2007. (1,2,3)Triazolo(4,5-h)quinolones. A new class of potent antitubercular agents against multidrug resistant *Mycobacterium tuberculosis* strains. *Bioorganic & Medicinal Chemistry Letters* 17(17), 4791-4794.
- Carta, A., Piras, S., Palomba, M., Jabes, D., Molicotti, P., Zanetti, S., 2008. Anti-Mycobacterial Activity of Quinolones. Triazoloquinolones a New Class of Potent Anti-Mycobacterial Agents. *Anti-Infective Agents in Med Chem* 7(2), 134-147.
- Carvalho, S.A., da Silva, E.F., de Souza, M.V.N., Lourenco, M.C.S., Vicente, F.R., 2008. Synthesis and antimycobacterial evaluation of new trans-cinnamic acid hydrazide derivatives. *Bioorganic & Medicinal Chemistry Letters* 18(2), 538-541.
- Chambhare, R.V., Khadse, B.G., Bobde, A.S., Bahekar, R.H., 2003. Synthesis and preliminary evaluation of some N-(5-(2-furanyl)-2-methyl-4-oxo-4H-thieno(2,3-d)pyrimidin-3-yl)-carboxamide and 3-substituted-5-(2-furanyl)-2-methyl-3H-thieno(2,3-d)pyrimidin-4-ones as antimicrobial agents. *European Journal of Medicinal Chemistry* 38(1), 89-100.
- Chhabria, M.T., Jani, M.H., 2009. Design, synthesis and antimycobacterial activity of some novel imidazo(1,2-c)pyrimidines. *European Journal of Medicinal Chemistry* 44(10), 3837-3844.
- Copp, B.R., Christiansen, H.C., Lindsay, B.S., Franzblau, S.G., 2005. Identification of heteroarylenamines as a new class of antituberculosis lead molecules. *Bioorganic & Medicinal Chemistry Letters* 15(18), 4097-4099.
- Das, U., Das, S., Bandy, B., Stables, J.P., Dimmock, J.R., 2008. N-Aroyl-3,5-bis(benzylidene)-4-piperidones: A novel class of antimycobacterial agents. *Bioorganic & Medicinal Chemistry* 16(7), 3602-3607.
- de Almeida, M.V., Saraiva, M.F., de Souza, M.V.N., da Costa, C.F., Vicente, F.R.C., Lourenco, M.C.S., 2007. Synthesis and antitubercular activity of lipophilic moxifloxacin and gatifloxacin derivatives. *Bioorganic & Medicinal Chemistry Letters* 17(20), 5661-5664.
- De Logu, A., Palchykovska, L.H., Kostina, V.H., et al., 2009. Novel N-aryl- and N-heteryl phenazine-1-carboxamides as potential agents for the treatment of infections sustained by drug-resistant and multidrug-resistant *Mycobacterium tuberculosis*. *International Journal of Antimicrobial Agents* 33(3), 223-229.
- de Souza, M.V.N., Pais, K.C., Kaiser, C.R., Peralta, M.A., Ferreira, M.L., Lourenço, M.C.S., 2009. Synthesis and in vitro antitubercular activity of a series of quinoline

- derivatives. *Bioorganic & Medicinal Chemistry* 17(4), 1474-1480.
- Desai, B., Sureja, D., Naliapara, Y., Shaha, A., Saxena, A.K., 2001. Synthesis and QSAR Studies of 4-Substituted Phenyl-2,6-dimethyl-3,5-Bis-N-(substituted Phenyl)carbamoyl-1,4-dihydropyridines as Potential Antitubercular Agents. *Bioorganic & Medicinal Chemistry* 9(8), 1993-1998.
- Devakaram, R.V., Yogeewari, P., China, A., Nagaraja, V., Sriram, D., 2008. Synthesis and antimycobacterial evaluation of newer 1-cyclopropyl-1,4-dihydro-6-fluoro-7-(substituted secondary amino)-8-methoxy-5-(sub)-4-oxoquinoline-3-carboxylic acids. *Bioorganic & Medicinal Chemistry* 16(5), 2558-2569.
- Dinakaran, M., Senthilkumar, P., Yogeewari, P., China, A., Nagaraja, V., Sriram, D., 2008. Antimycobacterial and phototoxic evaluation of novel 6-fluoro/nitro-4-oxo-7-(sub)-4H-(1,3)thiazeto(3,2-a)quinoline-3-carboxylic acid. *International Journal of Antimicrobial Agents* 31(4), 337-344.
- Dinakaran, M., Senthilkumar, P., Yogeewari, P., China, A., Nagaraja, V.M., Sriram, D., 2008. Antimycobacterial activities of novel 2-(sub)-3-fluoro/nitro-5, 12-dihydro-5-oxobenzo-thiazolo(3,2-a)quinoline-6-carboxylic acid. *Bioorganic & Medicinal Chemistry* 16(6), 3408-3418.
- Dinakaran, M., Senthilkumar, P., Yogeewari, P., China, A., Nagaraja, V., Sriram, D., 2008. Novel ofloxacin derivatives: Synthesis, antimycobacterial and toxicological evaluation. *Bioorganic & Medicinal Chemistry Letters* 18(3), 1229-1236.
- Dolezal, M., Cmedlova, P., Palek, L., et al., 2008. Synthesis and antimycobacterial evaluation of substituted pyrazinecarboxamides. *European Journal of Medicinal Chemistry* 43(5), 1105-1113.
- Drews, S.J., Hung, F., Av-Gay, Y., 2001. A protein kinase inhibitor as an antimycobacterial agent. *FEMS Microbiology Letter* 205(2), 369-374.
- Ducati, R.G., Ruffino-Netto, A., Basso, L.A., Santos, D.S., 2006. The resumption of consumption-A review on tuberculosis. *Mem Inst Oswaldo Cruz* 101(7), 697-714.
- El-Hamamsy, M.H.R.I., Smith, A.W., Thompson, A.S., Threadgill, M.D., 2007. Structurebased design, synthesis and preliminary evaluation of selective inhibitors of dihydrofolate reductase from *Mycobacterium tuberculosis*. *Bioorganic & Medicinal Chemistry* 15(13), 4552-4576.
- Eswaran, S., Adhikari, A.V., Pal, N.K., Chowdhury, I.H., 2010. Design and synthesis of some new quinoline-3-carbohydrazone derivatives as potential antimycobacterial agents. *Bioorganic & Medicinal Chemistry Letters* 20(3), 1040-1044.
- Fassihi, A., Azadpour, Z., Delbari, N., et al., 2009. Synthesis and antitubercular activity of novel 4-substituted imidazole-2,6-dimethyl-N3,N5-bisaryl-1,4-dihydropyridine-3,5-dicarboxamides. *European Journal of Medicinal Chemistry* 44(8), 3253-3258.
- Faugeroux, V., Genisson, Y., Salma, Y., Constant, P., Baltasa, M., 2007. Synthesis and biological evaluation of conformationally constrained analogues of the antitubercular agent ethambutol. *Bioorganic & Medicinal Chemistry* 15(17), 5866-5876.
- Figueiredo, R., Moiteiro, C., Medeiros, M.A., et al., 2009. Synthesis and evaluation of rifabutin analogs against *Mycobacterium avium* and H37Rv, MDR and NRP *Mycobacterium tuberculosis*. *Bioorganic & Medicinal Chemistry* 17(2), 503-511.
- Foks, H., Trapkowska, I., Janowiec, M., Zwolska, Z., Augustynowicz-Kopec, E., 2004. Studies on pyrazine derivatives. 38. Synthesis, reactions, and Tuberculostatic activity of pyrazinyl-substituted derivatives of hydrazinocarbodithioic acid. *Chem. Heterocyclic Compounds* 40(9), 1185-1193.
- Gaurrand, S., Desjardins, S., Meyer, C., et al., 2006. Conformational Analysis of R207910, a New Drug Candidate for the Treatment of Tuberculosis, by a Combined NMR and Molecular Modeling Approach. *Chem. Biol. Drug Des* 68(2), 77-84.
- Gaveriya, H., Desai, B., Vora, V., Shah, A., 2001. Synthesis of some new unsymmetrical 1,4-dihydropyridine derivatives as potent antitubercular agents. *Heterocyclic Communication* 5(5), 481-484.
- Gemma, S., Savini, L., Altarelli, M., et al., 2009. Development of antitubercular compounds based on a 4-quinolyhydrazone scaffold. Further structure-activity relationship studies. *Bioorganic & Medicinal Chemistry* 17(16), 6063-6072.
- Gezginci, M.H., Martin, A.R., Franzblau, S.G., 2001. Antimycobacterial Activity of Substituted Isosteres of Pyridine- and Pyrazinecarboxylic Acids. 2. *Journal of Medicinal Chemistry* 44(10), 1560-1563.
- Gratraud, P., Surolia, N., Besra, G.S., Surolia, A., Kremer, L., 2008. Antimycobacterial Activity and Mechanism of Action of NAS-91. *Antimicrob Agents Chemother* 52(3), 1162-1166.
- Gray, M.A., 1997. *Tuberculosis Drugs*. Orthopaedic Nursing, 16(4), 64-69.
- Guillon, J., Reynolds, R.C., Leger, J-M., et al., 2004. Synthesis and Preliminary In Vitro Evaluation of Antimycobacterial Activity of New Pyrrolo(1,2-a) quinoxalinecarboxylic Acid Hydrazone Derivatives. *Journal of Enzyme Inhibitor & Medicinal Chemistry* 19(6), 489-495.
- Gundersen, L.L., Nissen-Meyer, J., Spilsberg, B., 2002. Synthesis and antimycobacterial activity of 6-aryl purines: the requirement for the N-9 substituent in active

- antimycobacterial purines. *Journal of Medicinal Chemistry* 45(6), 1383-1386.
- Guo, S., Song, Y., Huang, Q., et al., 2010. Identification, Synthesis, and Pharmacological Evaluation of Tetrahydroindazole Based Ligands as Novel Antituberculosis Agents. *Journal of Medicinal Chemistry* 53(2), 649-659.
- He, X., Alian, A., de Montellano, P.R.O., 2007. Inhibition of the Mycobacterium tuberculosis enoyl acyl carrier protein reductase InhA by arylamides. *Bioorganic & Medicinal Chemistry* 15(21), 6649-6658.
- Hearn, M.J., Cynamon, M.H., Chen, M.F., et al., 2009. Preparation and antitubercular activities in vitro and in vivo of novel Schiff bases of isoniazid. *European Journal of Medicinal Chemistry* 44(10), 4169-4178.
- Hearn, M.J., Webster, E.R., Cynamon, M.H., 2005. Preparation and Properties of Antitubercular 1-Piperidino-3-Arylthioureas. *Journal of Heterocyclic Chemistry* 42(6), 1225-1229.
- Herzigova, P., Klimesovs, V., Palat, K., Kaustova, J., Dahse, H-M., Mollmann, U., 2009. Preparation and in-vitro Evaluation of 4-Benzylsulfanylpyridine-2-carbohydrazides as Potential Antituberculosis Agents. *Arch. Pharm. Chem. Life Sciences* 342(7), 394-404.
- Huang, Q., Mao, J., Wan, B., et al., 2009. Searching for New Cures for Tuberculosis: Design, Synthesis, and Biological Evaluation of 2-Methylbenzothiazoles. *Journal of Medicinal Chemistry* 52(21), 6757-6767.
- Hughes, M.A., McFadden, J.M., Townsend, C.A., 2005. New α -methylene- γ -butyrolactones with antimycobacterial properties. *Bioorganic & Medicinal Chemistry Letters* 15(17), 3857-3859.
- Hurdle, J.G., Lee, R.B., Budha, N.R., et al., 2008. A microbiological assessment of novel nitrofuranyl amides as anti-tuberculosis agents. *Journal of Antimicrob Chemother* 62(5), 1037-1045.
- Imramovsky, A., Polanc, S., Vinsova, J., et al., A new modification of anti-tubercular active molecules. *Bioorganic & Medicinal Chemistry* 15(7), 2551-2559.
- Jain, R., Vaitilingam, B., Nayyar, A., Palde, P.B., 2003. Substituted 4-Methylquinolines as a New Class of Anti-Tuberculosis Agents. *Bioorganic & Medicinal Chemistry Letters* 13(6), 1051-1054.
- Jaju, S., Palkar, M., Maddi, V., Ronad, P., Mamledesai, S., Satyanarayana, D., Ghatole, M., 2009. Synthesis and Antimycobacterial Activity of a Novel Series of Isonicotinylhydrazide Derivatives. *Arch. Pharm. Chem. Life Sciences* 342(12), 723-731.
- Janin, Y.L., 2007. Antituberculosis drugs: Ten years of research. *Bioorganic & Medicinal Chemistry* 15, 2479-2513 (2007).
- Japelj, B., Recnik, S., Cebašek, P., Stanovnik, B., Svete, J., 2005. Synthesis and Antimycobacterial Activity of Alkyl 1-Heteroaryl-1H-1,2,3-triazole-4-carboxylates. *Journal of Heterocyclic Chemistry* 42(6), 1167-1173.
- Jaso, A., Zarranz, B., Aldana, I., Monge, A., 2003. Synthesis of new 2-acetyl and 2-benzoyl quinoxaline 1,4-di-N-oxide derivatives as anti-Mycobacterium tuberculosis agents. *European Journal of Medicinal Chemistry* 38(9), 791-800.
- Jaso, A., Zarranz, B., Aldana, I., Monge, A., 2005. Synthesis of New Quinoxaline-2- carboxylate 1,4-Dioxide Derivatives as Anti-Mycobacterium tuberculosis Agents. *Journal of Medicinal Chemistry* 48(6), 2019-2025.
- Jean, L., Santos, J.L., Yamasaki, P.R., et al., 2009. Synthesis and in vitro anti Mycobacterium tuberculosis activity of a series of phthalimide derivatives. *Bioorganic & Medicinal Chemistry* 17(11), 3795-3799.
- Kamal, A., Ahmed, S.K., Reddy, K.S., et al., 2007. Anti-tubercular agents. Part IV: Synthesis and antimycobacterial evaluation of nitroheterocyclic-based 1,2,4-benzothiadiazines. *Bioorganic & Medicinal Chemistry Letters* 17(19), 5419-5422.
- Kamal, A., Babu, A.H., Ramana, A.V., Sinha, R., Yadav, J.S., Arora, S.K., 2005. Antitubercular agents. Part 1: Synthesis of phthalimido- and naphthalimido-linked phenazines as new prototype antitubercular agents. *Bioorganic & Medicinal Chemistry Letters* 15(7), 1923-1926.
- Kamal, A., Reddy, K.S., Ahmed, S.K., et al., 2006. Anti-tubercular agents. Part 3. Benzothiadiazine as a novel scaffold for anti-Mycobacterium activity. *Bioorganic & Medicinal Chemistry* 14(3), 650-658.
- Karthikeyan, S.V., Bala, B.D., Raja, V.P.A., Perumal, S., Yogeewari, P., Sriram, D., 2010. A highly atom economic, chemo-, regio- and stereoselective synthesis and evaluation of spiro-pyrrolothiazoles as antitubercular agents. *Bioorganic & Medicinal Chemistry Letters* 20(1), 350-353.
- Karthikeyan, S.V., Perumal, S., Shetty, K.A., Yogeewari, P., Sriram, D., 2009. A microwave-assisted facile regioselective Fischer indole synthesis and antitubercular evaluation of novel 2-aryl-3,4-dihydro-2H-thieno(3,2-b)indoles. *Bioorganic & Medicinal Chemistry Letters* 19(11), 3006-3009.
- Kawakami, K., Namba, K., Tanaka, M., Matsushashi, N., Sato, K., Takemura, M., 2000. Antimycobacterial Activities of Novel Levofloxacin Analogues. *Antimicrob Agents Chemother* 44(8), 2126-2129.
- Khan, A., Sarkar, S., Sarkar, D., 2008. Bactericidal activity of 2-nitroimidazole against the active replicating stage of Mycobacterium bovis BCG and Mycobacterium tuberculosis with intracellular efficacy in THP-1 macrophages. *International Journal of Antimicrobial Agents* 32(1), 40-45.

- Kharkar, P.S., Desai, B., Gaveria, H., et al., 2002. Three dimensional quantitative structureactivity relationship of 1,4-dihydropyridines as antitubercular agents. *Journal of Medicinal Chemistry* 45(22), 4858-4867.
- Khoshneviszadeh, M., Edraki, N., Javidnia, K., et al., 2009. Synthesis and biological evaluation of some new 1,4-dihydropyridines containing different ester substitute and diethyl carbamoyl group as anti-tubercular agents. *Bioorganic & Medicinal Chemistry* 17(4), 1579-1586.
- Kim, P., Kang, S., Boshoff, H.I., et al., 2009. Structure-Activity Relationships of Antitubercular Nitroimidazoles. 2. Determinants of Aerobic Activity and Quantitative Structure-Activity Relationships. *Journal of Medicinal Chemistry* 52(5), 1329-1344.
- Kim, P., Zhang, L., Manjunatha, U.H., et al., 2009. Structure-Activity Relationships of Antitubercular Nitroimidazoles. 1. Structural Features Associated with Aerobic and Anaerobic Activities of 4- and 5-Nitroimidazoles. *Journal of Medicinal Chemistry* 52(5), 1317-1328.
- Kini, S.G., Bhat, A.R., Bryant, B., Williamson, J.S, Dayan, F.E., 2009. Synthesis, antitubercular activity and docking study of novel cyclic azole substituted diphenyl ether derivatives. *European Journal of Medicinal Chemistry* 44(2), 492-500.
- Koketsu, M., Tanaka, K., Takenaka, Y., Kwong, C.D., Ishihara, H., 2002. Synthesis of 1,3 thiazine derivatives and their evaluation as potential antimycobacterial agents. *European Journal of Pharmaceutical Sciences* 15(3), 307-310.
- Krinkova, J., Dolezal, M., Hartl, J.V, Buchta, V., Pour, M., 2002. Synthesis and biological activity of 5-alkyl-6-alkylsulfanyl or 5-alkyl-6-arylsulfanyl pyrazine-2-carboxamides and corresponding thioamides. *II Farmaco* 57(1), 71-78.
- Kumar, A., Sinha, S., Chauhan, P.M.S., 2002. Synthesis of novel antimycobacterial combinatorial libraries of structurally diverse substituted pyrimidines by three-Component solid-phase reactions. *Bioorganic & Medicinal Chemistry Letters* 12(4), 667-669.
- Kumar, R.R., Perumal, S., Senthilkumar, P., Yogeewari, P., Sriram, D., 2007. An atom efficient, solvent-free, green synthesis and antimycobacterial evaluation of 2-amino-6-methyl-4-aryl-8-((E)-arylmethylidene)-5,6,7,8-tetrahydro-4Hpyrano(3,2-c)pyridine-3-carbonitriles. *Bioorganic & Medicinal Chemistry Letters* 17(23), 6459-6462.
- Kumar, R.R., Perumal, S., Senthilkumar, P., Yogeewari, P., Sriram, D., 2008. Discovery of Antimycobacterial Spiropiperidin-4-ones: An Atom Economic, Stereoselective Synthesis, and Biological Intervention. *Journal of Medicinal Chemistry* 51(18), 5731-5735.
- Li, X., Manjunatha, U.H., Goodwin, M.B., et al., 2008. Synthesis and antitubercular activity of 7-(R)- and 7-(S)-methyl-2-nitro-6-(S)-(4-(trifluoromethoxy)benzyloxy)-6,7-dihydro-5H-imidazo(2,1-b)(1,3)oxazines, analogues of PA-824. *Bioorganic & Medicinal Chemistry Letters* 18(7), 2256-2262.
- Lilienkampf, A., Mao, J., Wan, B., Wang, Y., Franzblau, S.G., Kozikowski, A.P., 2009. Structure-Activity Relationships for a Series of Quinoline-Based Compounds Active against Replicating and Nonreplicating Mycobacterium tuberculosis. *Journal of Medicinal Chemistry* 52(7), 2109-2118.
- Lin, Y-M., Zhou, Y., Flavin, M.T., Zhou, L-M., Nie, W., Chen, F-C., 2002. Chalcones and Flavonoids as Anti-Tuberculosis Agents. *Bioorganic & Medicinal Chemistry* 10(8), 2795-2802.
- Loddenkemper, R., Sagebiel, D., Brendel, A., 2002. Strategies against multidrug-resistant tuberculosis. *European Respir Journal* 20(36), 66s-77s.
- Lourenco, M.C.S., Ferreira, M.L., de Souza, M.V.N., Peralta, M.A., Vasconcelos, T.R.A., Henriques, M.G.M.O., 2008. Synthesis and anti-mycobacterial activity of (E)-N'-(monosubstituted-benzylidene)isonicotinohydrazide derivatives. *European Journal of Medicinal Chemistry* 43(6), 1344-1347.
- Maccari, R., Ottana, R., Monforte, F., Vigorita, M.G., 2002, In *Vitro* Antimycobacterial Activities of 2'-Monosubstituted Isonicotinohydrazides and Their Cyanoborane Adducts. *Antimicrob Agents Chemother* 46(2), 294-299.
- Madrid, P.B., Polgar, W.E., Toll, L., Tangaa, M.J., 2007. Synthesis and antitubercular activity of phenothiazines with reduced binding to dopamine and serotonin receptors. *Bioorganic & Medicinal Chemistry Letters* 17(11), 3014-3017.
- Makarov, V., Manina, G., Mikusova, K., et al., 2009. Benzothiazinones Kill *Mycobacterium tuberculosis* by Blocking Arabinan Synthesis. *Science* 324(5928), 801-804.
- Mamolo, M.G., Zampier, D., Falagiani, V., Vio, L., Banfi, E., 2001. Synthesis and antimycobacterial activity of 5-aryl-1-isonicotinoyl-3-(pyridin-2-yl)-4,5-dihydro-1H-pyrazole derivatives. *II Farmaco* 56(8), 593-599.
- Mao, J., Yuan, H., Wang, Y., et al., 2009. From Serendipity to Rational Antituberculosis Drug Discovery of Mefloquine-Isoxazole Carboxylic Acid Esters. *Journal of Medicinal Chemistry* 52(22), 6966-6978.
- Matlola, N.M., Steel, H.C., Anderson, R., 2001. Antimycobacterial action of B4128, a novel tetramethylpiperidyl-substituted phenazine. *Journal of Antimicrob. Chemother* 47(2), 199-202.
- Monga, V., Nayyar, A., Vaitilingam, B., et al., 2004. Ring-substituted quinolines. Part 2: Synthesis and antimycobacterial activities of ring-substituted quinolinecarbohydrazide and ring-substituted quinolinecarboxamide analogues. *Bioorganic & Medicinal Chemistry* 12(24), 6465-6472.

- Morgan, J., Haritakul, R., Keller, P.A., 2003. Anilino pyrimidines as novel antituberculosis agents. *Bioorganic & Medicinal Chemistry Letters* 13(10), 1755-1757.
- Moukha-Chafiq, O., Taha, M.L., Lazrek, H.B., et al., 2001. Synthesis and biological evaluation Of some 4-substituted 1-(1-(4-hydroxybutyl)-1,2,3-triazol-(4&5)-ylmethyl)-1H-pyrazolo-(3,4-d)pyrimidines. *Nucleosides, Nucleotides & Nucleic acids* 20(10&11), 1811-1821.
- Moukha-Chafiq, O., Taha, M.L., Lazrek, H.B., Vasseur, J.J., Clercq, E.D., 2002, Synthesis and biological evaluation of some acyclic a-(1H-pyrazolo-(3,4-d)pyrimidin-4-yl)thioalkylamide Nucleosides. *Nucleosides, Nucleotides and Nucleic acids*, 21(2), 165-176.
- Moukha-Chafiq, O., Taha, M.L., Mouna, A., 2007. Synthesis and biological evaluation of some A-(6-(1/-carbamoylethylthio)-1H-pyrazolo(3,4-d)pyrimidin-4-yl)thioalkylcarboxamide Acyclonucleosides. *Nucleosides, Nucleotides and Nucleic acids*, 26(4), 335-345.
- Nagaraj, A., Reddy, C.S., 2007. Synthesis and Biological Study of Novel Methylene-bis(chalcones and Substituted Methylene-bis-pyrimidines/pyrimidinones. *Journal of Heterocyclic Chemistry* 44(5), 1181-1185.
- Nayyar, A., Patel, S.R., Shaikh, M., Coutinho, E., Jain, R., 2009. Synthesis, anti-tuberculosis activity and 3D-QSAR study of amino acid conjugates of 4-(adamantan-1-yl) group containing quinolines. *European Journal of Medicinal Chemistry* 44(5), 2017-2029.
- Odell, L.R., Nilsson, M.T., Gising, J., et al., 2009. Functionalized 3-amino-imidazo(1,2-a)pyridines: A novel class of drug-like Mycobacterium tuberculosis glutamine synthetase inhibitors. *Bioorganic & Medicinal Chemistry Letters* 19(16), 4790-4793.
- Onajole, O.K., Govender, K., Govender, P., et al., 2009. Pentacyclo-undecane derived cyclic tetra-amines: Synthesis and evaluation as potent anti-tuberculosis agents. *European Journal of Medicinal Chemistry* 44(11), 4297-4305.
- Opletalova, V., Hartl, J., Patel, A., Palat, Jr K., Buchta, V., 2002. Ring substituted 3-phenyl- 1-(2 pyrazinyl)-2 propen-1-ones as potential photosynthesis inhibiting, antifungal and antimycobacterial agents. *II Farmaco* 57(2), 135-144.
- Ozdemir, A., Turan-Zitouni, G., Kaplancikli, Z.A., Tunali, Y., 2009. Synthesis and biological activities of new hydrazide derivatives. *Journal of Enzyme Inhibitor & Medicinal Chemistry* 24(3), 825-831.
- Pagani, G., Pregnolato, M., Ubiali, D., et al., 2000. Synthesis and in Vitro Anti- Mycobacterium Activity of N-Alkyl-1,2-dihydro-2-thioxo-3-pyridinecarbothioamides. Preliminary Toxicity and Pharmacokinetic Evaluation. *Journal of Medicinal Chemistry* 43(2), 199-204.
- Panda, G., Parai, M.K., Das, S.K., et al., 2007. Effect of substituents on diarylmethanes for antitubercular activity. *European Journal of Medicinal Chemistry* 42(3), 410-419.
- Parai, M.K., Panda, G., Chaturvedi, V., Manju, Y.K., Sinha, S., 2008. Thiophene containing triarylmethanes as antitubercular agents. *Bioorganic & Medicinal Chemistry Letters* 18(1), 289- 292.
- Pathak, A.K., Pathak, V., Seitz, L.E., Suling, W.J., Reynolds, R.C., 2004. Antimycobacterial Agents. 1. Thio Analogues of Purine. *Journal of Medicinal Chemistry* 47(1), 273-276.
- Perri, G.D., Bonora, S., 2004. Which agents should we use for the treatment of multidrugresistant Mycobacterium tuberculosis?. *Journal of Antimicrob Chemother* 54(3), 593-602.
- Pironi, M., Lilienkampf, A., Wan, B., Wang, Y., Franzblau, S.G., Kozikowski, A.P., 2009. Synthesis, Biological Evaluation, and Structure-Activity Relationships for 5-((E)-2-Arylethenyl)-3-isoxazolecarboxylic Acid Alkyl Ester Derivatives as Valuable Antitubercular Chemotypes. *Journal of Medicinal Chemistry* 52(20), 6287-6296.
- Ragno, R., Marshall, G.R., Santo, R.D., et al., 2000. Antimycobacterial Pyrroles: Synthesis, Anti-Mycobacterium tuberculosis Activity and QSAR Studies. *Bioorganic & Medicinal Chemistry* 8(6), 1423-1432.
- Rakesh., Sun, D., Lee, R.B., Tangallapally, R.P., Lee, R.E., 2009. Synthesis, optimization and structureactivity relationships of 3,5-disubstituted isoxazolines as new antituberculosis agents. *European Journal of Medicinal Chemistry* 44(2), 460-472.
- Raparti, V., Chitre, T., Bothara, K., et al., 2009. Novel 4-(morpholin-4-yl)-N'-(arylidene) benzohydrazides: Synthesis, antimycobacterial activity and QSAR investigations. *European Journal of Medicinal Chemistry* 44(10), 3954-3960.
- Reddy, V.M., O'Sullivan, J.F., Gangadharam, P.R., 1999. Antimycobacterial activities of riminophenazines. *Journal of Antimicrob Chemother* 43(5), 615-623.
- Savini, L., Chiasserini, L., Gaeta, A., Pellerano, C., 2002. Synthesis and Anti-tubercular Evaluation of 4-Quinolylhydrazones. *Bioorganic & Medicinal Chemistry* 10(7), 2193-2198.
- Scozzafava, A., Mastrolorenzo, A., Supuran, C.T., 2001. Antimycobacterial activity of 9- sulphonylated/ sulphenylated-6-mercaptapurine derivatives. *Bioorganic & Medicinal Chemistry Letters* 11(13), 1675-1678.
- Scozzafava, A., Mastrolorenzo, A., Supuran, C.T., 2001. Antimycobacterial Activity of 3,4-dichlorophenyl-ureas, N,N-diphenyl-ureas and Related Derivatives. *Journal of Enzyme Inhibitors* 16(3), 425-432.
- Seitz, L.E., Suling, W.J., Reynolds, R.C., 2002. Synthesis and antimycobacterial activity of pyrazine and quinoxaline

- derivatives. *Journal of Medicinal Chemistry* 45(25), 5604-5606.
- Senthilkumar, P., Dinakaran, M., Chandrasekaran, Y., Yogeewari, P., Sriram, D., 2009. Synthesis and in-vitro Antimycobacterial Evaluation of 1-(Cyclopropyl/2,4-difluorophenyl/tert-butyl)-1,4-dihydro-8-methyl-6-nitro-4-oxo-7-(substituted secondary amino)quinoline-3-carboxylic acids. *Arch Pharm Chem Life Sciences* 342(2), 100-112.
- Senthilkumar, P., Dinakaran, M., Yogeewari, P., China, A., Nagaraja, V., Sriram, D., 2009. Antimycobacterial activities of novel fluoroquinolones. *Biomedicine Pharmacother.* 63(1), 27-35.
- Senthilkumar, P., Dinakaran, M., Yogeewari, P., China, A., Nagaraja, V., Sriram, D., 2009. Antimycobacterial activities of novel fluoroquinolones. *Biomedicine Pharmacother.* 63(1), 27-35.
- Senthilkumar, P., Dinakaran, M., Yogeewari, P., Sriram, D., China, A., Nagaraja, V., 2009. Synthesis and antimycobacterial activities of novel 6-nitroquinolone-3-carboxylic acids. *European Journal of Medicinal Chemistry* 44(1), 345-358.
- Sharma, M., Chaturvedi, V., Manju, Y.K., et al., 2009. Substituted quinolinyl chalcones and quinolinyl pyrimidines as a new class of anti-infective agents. *European Journal of Medicinal Chemistry* 44(5), 2081-2091.
- Sheu, J-Y., Chen, Y-L., Tzeng, C-C., Hsu S-L., Fang, K-C., Wang, T-C., 2003. Synthesis, and Antimycobacterial and Cytotoxic Evaluation of Certain Fluoroquinolone Derivatives. *Helv Chimica Acta* 86(7), 2481-2489.
- Silva, R.S.F., Pinto, M.C.F.R., Goulart, M.O.F., et al., 2009. A macrolactone from benzo(a)phenazine with potent activity against *Mycobacterium tuberculosis*. *European Journal of Medicinal Chemistry* 44(5), 2334-2337.
- Sriram, D., Aubry, A., Yogeewari, P., Fisher, L.M., 2006. Gatifloxacin derivatives: Synthesis, antimycobacterial activities, and inhibition of *Mycobacterium tuberculosis* DNA gyrase. *Bioorganic & Medicinal Chemistry Letters* 16(11), 2982-2985.
- Sriram, D., Yogeewari, P., Basha, J.S., Radha, D.R., Nagaraja, V., 2005. Synthesis and antimycobacterial evaluation of various 7-substituted ciprofloxacin derivatives. *Bioorganic & Medicinal Chemistry* 13(20), 5774-5778.
- Sriram, D., Yogeewari, P., Devakaram, R.V., 2006. Synthesis, in vitro and in vivo antimycobacterial activities of diclofenac acid hydrazones and amides. *Bioorganic & Medicinal Chemistry* 14(9), 3113-3118.
- Sriram, D., Yogeewari, P., Dhakla, P., Senthilkumar, P., Banerjee, D., Manjashetty, T.H., 2009. 5-Nitrofuran-2-yl derivatives: Synthesis and inhibitory activities against growing and dormant mycobacterium species. *Bioorganic & Medicinal Chemistry Letters* 19(4), 1152-1154.
- Sriram, D., Yogeewari, P., Dinakaran, M., Thirumurugan, R., 2007. Antimycobacterial activity of novel 1-(5-cyclobutyl-1,3-oxazol-2-yl)-3-(sub)phenyl/pyridylthiourea compounds endowed with high activity toward multidrug-resistant *Mycobacterium tuberculosis*. *Journal of Antimicrob Chemother* 59(6), 1194-1196.
- Sriram, D., Yogeewari, P., Gopal, G., 2005. Synthesis, anti-HIV and antitubercular activities of lamivudine prodrugs. *European Journal of Medicinal Chemistry* 40(12), 1373-1376.
- Sriram, D., Yogeewari, P., Madhu, K., 2005. Synthesis and in vitro and in vivo antimycobacterial activity of isonicotinoyl hydrazones. *Bioorganic & Medicinal Chemistry Letters* 15(20), 4502-4505.
- Sriram, D., Yogeewari, P., Madhu, K., 2006. Synthesis and in vitro antitubercular activity of some 1-((4-sub)phenyl)-3-(4-{1-((pyridine-4-carbonyl) hydrazono)ethyl}phenyl)thiourea. *Bioorganic & Medicinal Chemistry Letters* 16(4), 876-878.
- Sriram, D., Yogeewari, P., Senchani, G., Banerjee, D., 2007. Newer tetracycline derivatives: Synthesis, anti-HIV, antimycobacterial activities and inhibition of HIV-1 integrase. *Bioorganic & Medicinal Chemistry Letters* 17(8), 2372-2375.
- Sriram, D., Yogeewari, P., Thirumurugan, R., Pavana, R.K., 2006. Discovery of New Antitubercular Oxazolyl Thiosemicarbazones. *Journal of Medicinal Chemistry* 49(12), 3448-3450.
- Sun, D., Scherman, M.S., Jones, V., et al., 2009. Discovery, synthesis, and biological evaluation of piperidinol analogs with anti-tuberculosis activity. *Bioorganic & Medicinal Chemistry* 17(10), 3588-3594.
- Sunduru, N., Sharma, M., Chauhan, P.M.S., 2010. Recent advances in the design and synthesis of Heterocycles as anti-tubercular agents. *Future Medicinal Chemistry* 2(9), 1469-1500.
- Talatha, S., Gadad, A.K., 2006. Synthesis, antibacterial and antitubercular activities of some 7-(4-(5-amino-(1,3,4)thiadiazole-2-sulfonyl)-piperazin-1-yl) fluoroquinolonic derivatives. *European Journal of Medicinal Chemistry* 41(8), 918-924.
- Tangallapally, R.P., Lee, R.B., Lenaerts, A.J., Lee, R.E., 2006. Synthesis of new and potent analogues of anti-tuberculosis agent 5-nitro-furan-2-carboxylic acid 4-(4-benzylpiperazin-1-yl)-benzylamide with improved bioavailability. *Bioorganic & Medicinal Chemistry Letters* 16(10), 2584-2589.
- Tangallapally, R.P., Sun, D., Rakesh, et al., 2007. Discovery of novel isoxazolines as antituberculosis agents. *Bioorganic & Medicinal Chemistry Letters* 17(23), 6638-6642.
- Tangallapally, R.P., Yendapally, R., Daniels, A.J., Lee, R.B., Lee, R.E., Nitrofurans as novel anti-tuberculosis agents:

- identification, development and evaluation. *Current Topic in Medicinal Chemistry* 7(5), 509-526 (2007).
- Tangallapally, R.P., Yendapally, R., Lee, R.E., Lenaerts, A.J.M., Lee, R.E., 2005. Synthesis and evaluation of cyclic secondary amine substituted phenyl and benzyl nitrofuranyl amides as novel antituberculosis agents. *Journal of Medicinal Chemistry* 48(26), 8261-8269.
- Thompson, A.M., Blaser, A., Anderson, R.F., et al., 2009. Synthesis, Reduction Potentials, and Antitubercular Activity of Ring A/B Analogues of the Bioreductive Drug (6S)-2-Nitro-6-((4-(trifluoromethoxy)benzyl)oxy)-6,7-dihydro-5H-imidazo(2,1-b)(1,3)oxazine (PA-824). *Journal of Medicinal Chemistry* 52(3), 637-645.
- Tripathi, R.P., Verma, S.S., Pandey, J., et al., 2006. Search of antitubercular activities in tetrahydroacridines: Synthesis and biological evaluation. *Bioorganic & Medicinal Chemistry Letters* 16(19), 5144-5147.
- Ukrainets, I.V., Mospanova, E.V., Sidorenko, L.V., 2007. 4-hydroxy-2-quinolones. 1-hydroxy-3-oxo-5,6-dihydro-3h-pyrrolo(3,2,1-ij)-Quinoline-2-carboxylic acid hetaryl amides As potential antitubercular agents. *Chemistry of Heterocyclic Compounds* 43(7), 863-870.
- Upadhayaya, R.S., Kulkarni, G.M., Vasireddy, N.R., et al., 2009. Design, synthesis and biological evaluation of novel triazole, urea and thiourea derivatives of quinoline against *Mycobacterium tuberculosis*. *Bioorganic & Medicinal Chemistry* 17(13), 4681-4692.
- Upadhayaya, R.S., Vandavasi, J.K., Vasireddy, N.R., Sharma, V., Dixit, S.S., Chattopadhyaya, J., 2009. Design, synthesis, biological evaluation and molecular modelling studies of novel quinoline derivatives against *Mycobacterium tuberculosis*. *Bioorganic & Medicinal Chemistry* 17(11), 2830-2841.
- Velezheva, V.S., Brennan, P.J., Marshakov, V.Y., et al., 2004. Novel Pyridazino(4,3-b)indoles with Dual Inhibitory Activity against *Mycobacterium tuberculosis* and Monoamine Oxidase. *Journal of Medicinal Chemistry* 47(13), 3455-3461.
- Velikorodov, A.V., Urlyapova, N.G., Daudova, A.D., 2006. Synthesis and antimycobacterial activity of carbamate derivatives of 1,2-oxazine. *Pharmaceutical Chemistry Journal* 40(7), 380-382.
- Vicente, E., Perez-Silanes, S., Lima, L.M., et al., 2009. Selective activity against *Mycobacterium tuberculosis* of new quinoxaline 1,4-di-N-oxides. *Bioorganic & Medicinal Chemistry* 17(1), 385-389.
- Villar, R., Vicente, E., Solano, B., et al., 2008. In vitro and in vivo antimycobacterial activities of ketone and amide derivatives of quinoxaline 1,4-di-N-oxide. *Journal of Antimicrob Chemother* 62(3), 547-554.
- Virsoadia, V., Pissurlenkar, R.R.S., Manvar, D., et al., Synthesis, screening for antitubercular activity and 3D-QSAR studies of substituted N-phenyl-6-methyl-2-oxo-4-phenyl-1,2,3,4-tetrahydro-pyrimidine-5-carboxamides. *European Journal of Medicinal Chemistry* 43(10), 2103-2115 (2008).
- Waisser, K., Dražková, K., Kuneš, J., Klimešová, V., Kaustová, J., 2004. Antimycobacterial N-pyridinylsalicylamides, isosters of salicylamides. *II Farmaco* 59(8), 615-625.
- Weis, R., Schweiger, K., Faist, J., et al., 2008. Antimycobacterial and H1-antihistaminic activity of 2-substituted piperidine derivatives. *Bioorganic & Medicinal Chemistry* 16(24), 10326-10331.
- Weis, R., Faist, J., di Vora, U., et al., 2008. Antimycobacterial activity of diphenylpyraline derivatives. *European Journal of Medicinal Chemistry* 43(4), 872-879.
- Zarranz, B., Jaso, A., Aldana, I., Monge, A., 2003. Synthesis and Antimycobacterial Activity of New Quinoxaline-2-carboxamide 1,4-di-N-Oxide Derivatives. *Bioorganic & Medicinal Chemistry* 11(10), 2149-2156.
- Zhang, X., Hu, Y., Chen, S., et al., 2009. Synthesis and evaluation of (S,S)-N,N'-bis-(3-(2,2',6,6'-tetramethylbenzhydryloxy)-2-hydroxy-propyl)-ethylenediamine (S2824) analogs with anti-tuberculosis activity. *Bioorganic & Medicinal Chemistry Letters* 19(21), 6074-6077.
- Zhang, Y., Jin, G., Illarionov, B., Bacher, A., Fischer, M., Cushman, M., 2007. A New Series of 3-Alkyl Phosphate Derivatives of 4,5,6,7-Tetrahydro-1-D-ribityl-1H-pyrazolo(3,4-d) pyrimidine-dione as Inhibitors of Lumazine Synthase: Design, Synthesis, and Evaluation. *Journal of Organic Chemistry* 72(19), 7176-7184.

Visit us at: <http://bosajournals.com/chemint/>

Submissions are accepted at: editorci@bosajournals.com