



## Kinetic and thermodynamic properties of pharmaceutical drug (Gabapentin) by potassium bromate ( $\text{KBrO}_3$ ) in presence of micro amount of Ir(III) chloride as catalyst in acidic medium

Reena Patel<sup>1</sup>, Shailesh kumar<sup>1</sup>, Abhishek Verma<sup>1</sup> and Sheila Srivastava<sup>2,\*</sup>

<sup>1</sup>Department of Applied Chemistry B. B. A. U. Lucknow U.P., India

<sup>2</sup>Department of Chemistry Feroze Gandhi College, Raebareli U.P., India

\*Corresponding author's E. mail: [she\\_ila72@yahoo.com](mailto:she_ila72@yahoo.com), [patel.reena14@gmail.com](mailto:patel.reena14@gmail.com)

### ARTICLE INFO

#### Article type:

Research article

#### Article history:

Received March 2016

Accepted July 2016

April 2017 Issue

#### Keywords:

Kinetics

Ir(III) chloride

Oxidation

Gabapentin

Potassium bromate

Acidic medium

### ABSTRACT

The kinetic and thermodynamic properties of pharmaceutical drug (gabapentine) by potassium bromate ( $\text{KBrO}_3$ ) in presence of Micro amount of Ir(III) chloride as catalyst in acidic medium was studied in the temperature range 30 to 45<sup>o</sup> C. The reaction is carried out in the presence of mercuric acetate as a scavenger for chloride ion. 1-carboxy cyclohexane l-acetic acid was obtained as the oxidation product and identified chromatographically. The rate law followed a first order and zero order dependence with respect to  $\text{KBrO}_3$  and GBP respectively. The reaction followed first order with respect to Ir(III) chloride and  $[\text{H}^+]$ . Negligible effect of  $[\text{Hg}(\text{OAc})_2]$  and ionic strength of the medium was observed. Chloride ion positively influenced the rate of reaction. The values of rate constants observed at different temperatures (30 to 45<sup>o</sup>C) were utilized to calculate the activation parameters. Feasible mechanism is proposed which are be composed with the kinetics, stoichiometry and product of the reaction. The rate law has been derived from obtained kinetic data.

© 2017 International Scientific Organization: All rights reserved.


**Capsule Summary:** Effect of Ir(III) chloride and temperature on kinetic and thermodynamic properties of gabapentine was studied and study revealed that both catalyst and temperature affected both kinetic and thermodynamic properties considerably.

**Cite This Article As:** Reena Patel, Shailesh kumar, Abhishek Verma and Sheila Srivastava. Kinetic and thermodynamic properties of pharmaceutical drug (Gabapentin) by potassium bromate ( $\text{KBrO}_3$ ) in presence of micro amount of Ir(III) chloride as catalyst in acidic medium. Chemistry International 3(2) (2017) 158-164.

### INTRODUCTION

The transition metal catalysed reactions are important for the chemical industry from both practical and economic point of view. Transition metal ions are found to be good catalysts and their complexes are also able to catalyze a wide variety of reactions like hydrogenation, oxidation and

polymerization. The applications of transition metal catalyst such as Ru(III) (Singh et al., 2007; Hosahalli et al., 2012), Rh(III) (Singh et al., 2009 and 2010), Cu(II) (Li et al., 2009; Olusanya and Odebunmi, 2011) and Pd(II) (Koli and Nandibewoor, 2000; Singh et al., 2010) in kinetic studies of redox reaction involving organic substrate are reported in literature. It was found that these catalysts work efficiently in both acidic and alkaline media. The use of Ir(III) chloride as a



**Scheme 1:** Stoichiometry of Ir(III) catalyzed oxidation of gabapentin by potassium bromate [KBrO<sub>3</sub>]

non-toxic and homogenous catalyst has been reported (Bai et al., 2008; Munavalli et al., 2008; Singh et al., 2004; Wen-yu et al., 2005). The reaction path depends not only upon the nature of the oxidant and the nature of the substrates but also upon the ways in which reactive species of transition metal ion forms complex with the reactant molecules before changing into final products under experimental conditions. Oxidant such as NBS (Goel and Sharma, 2012; Mohana K N & Ramya, 2009), NBA (Singh et al., 2007), Periodate (Kaushik et al., 2010; Chaturvedi and Mishra, 2010), and iodate (Chanakira et al., 2006; Singh et al., 2011), have been successfully utilized in kinetic and mechanistic investigations of various organic substrate. Amongst various inorganic oxidants, iodate has been used as an oxidant in the uncatalyzed oxidation of oxalic acid (Ramana and Rao, 1991), acetophenone (Munikyamba et al., 1983), benzaldehyde (Munikyamba et al., 1981). Others introduced various reports in the literature regarding the oxidation of reducing sugars involving Ir(III) as homogenous catalyst (Brahmaiah and Manikyamba, 1995; Muthakia and Jonnalagadda, 1989; Mambo and Simoyi, 1993; Radhakrishnamurti and Tripathy, 1986).

Potassium bromate is known to be a powerful oxidizing agent with redox potential 1.44 volts and has been used widely in the oxidation of several organic substrates (Radhakrishnamurthy and Sarangi, 1981). However, till date no report is available in which bromate as an oxidant and Ir(III) as homogeneous catalyst have been used in the oxidation of gabapentin in acidic medium. Gabapentin (GBP) is most potent drug prescribed usually in combination with other medications for the prevention of seizure. It is sometimes prescribed for the management of neuralgia (nerve pain) (Jensen et al., 2002). Gabapentin has been prescribed for the treatment of some mood disorders, anxiety etc. The study of neuroleptic drug is more important due to biological significance of these drugs and their selectivity towards the oxidation to yield different products.

## MATERIAL AND METHODS

### Material


Aqueous solution of Gabapentin (CDH), potassium bromate (S.D. Fine A.R.) and mercuric acetate (E. Merck) were prepared by dissolving the weighed amount of sample in triple distilled water. Perchloric acid (60% E. Merck) was used as a source of hydrogen ions. Iridium tri chloride (Johnson Matthey) was prepared by dissolving the sample in hydrochloric acid of known strength. All other reagents of analytical grade were available. Sodium perchlorate (E. Merck) was used to maintain the ionic strength of the medium. The reaction stills were blackened from outside to prevent photochemical effect.

### Kinetic Procedure

Aliquots (5ml) of the reaction mixture were pipetted out at regular intervals of time and poured in to a conical flask containing 5 ml of 4 % KI solution and 5 ml of dilute sulfuric acid. The liberated bromine equivalent to consumed oxidant was estimated with standard sodium thiosulphate solution using starch as an indicator. The initial rates were obtained from slopes of concentration vs. time graph in the initial stages of the reaction by plane mirror method.

### Stoichiometry of Ir(III) catalyzed oxidation of gabapentin by potassium bromate [KBrO<sub>3</sub>]

The stoichiometry of the reaction was determined by equilibrating varying ratios of KBrO<sub>3</sub> and [GBP] at 35°C for 48 hours under kinetic condition. Estimation of unconsumed KBrO<sub>3</sub> revealed that, one mole of gabapentin consumes 2 moles of potassium bromate. This result confirms 1:2 stoichiometry. According to the above reaction the oxidation product of gabapentin is 1-carboxycyclohexane-1-acetic acid, which was identified and confirmed by paper chromatography. 1-carboxycyclohexane-1-acetic acid was identified and confirmed by IR spectral analysis. The bands at 1760 cm<sup>-1</sup> and 1710 cm<sup>-1</sup> corresponds to two-C=O group and 3550 cm<sup>-1</sup> corresponds to two -OH group clearly confirms 1-carboxycyclohexane-1-acetic acid (Scheme 1).



Scheme 2: Reaction mechanism

**Table 1:** Effect of variation of oxidant, substrate and catalyst at 35°C

Oxidant x 10 <sup>3</sup> M (Potassium bromate)	[Substrate]x 10 <sup>2</sup> M (Gabapentin)	Ir(III) x 10 <sup>5</sup> M	(-dc/dt)x10 <sup>7</sup> ML <sup>-1</sup> s <sup>-1</sup>
0.80	1.00	1.00	1.30
1.00	1.00	1.00	2.00
1.25	1.00	1.00	2.10
1.69	1.00	1.00	3.20
2.50	1.00	1.00	6.20
5.00	1.00	1.00	13.00
1.00	0.33	1.00	2.21
1.00	0.44	1.00	2.00
1.00	0.55	1.00	2.10
1.00	0.66	1.00	2.31
1.00	1.00	1.00	2.11
1.00	2.00	1.00	2.22
1.00	1.00	0.66	1.00
1.00	1.00	1.33	2.00
1.00	1.00	2.67	2.62
1.00	1.00	4.01	3.50
1.00	1.00	5.34	6.00
1.00	1.00	6.67	11.3


**Table 2:** Effect of variation of reactant (HClO<sub>4</sub> and KCl, NaClO<sub>4</sub>) at 35°C

HClO <sub>4</sub> x 10 <sup>3</sup> M	[KCl] x 10 <sup>3</sup> M	Hg(OAc) <sub>2</sub> x 10 <sup>3</sup> M	(-dc/dt)x10 <sup>7</sup> ML <sup>-1</sup> s <sup>-1</sup>
0.80	1.00	1.00	1.20
1.00	1.00	1.00	2.00
1.25	1.00	1.00	2.30
1.69	1.00	1.00	4.20
2.50	1.00	1.00	6.20
5.00	1.00	1.00	12.00
1.00	0.80	1.00	1.10
1.00	1.00	1.00	2.00
1.00	1.25	1.00	2.80
1.00	1.69	1.00	3.70
1.00	2.50	1.00	5.20
1.00	5.00	1.00	9.31
1.00	1.00	0.80	2.00
1.00	1.00	1.00	2.50
1.00	1.00	1.25	2.22
1.00	1.00	1.69	3.00
1.00	1.00	2.50	2.00
1.00	1.00	5.00	2.13


## RESULTS AND DISCUSSION

The kinetic results were collected at several initial concentrations of reactants (Table 1). First order rate constant  $k_1$  i.e.  $(-dc/dt/KBrO_3^*)$  were calculated from the


plots of unconsumed potassium bromate vs. time. The plots of  $\log(-dc/dt)$  versus  $\log(\text{oxidant})$  were linear indicating first order dependence on  $KBrO_3$  (Fig-1). Insignificant effect on the rate was observed on increasing the concentration of



**Fig. 1:** Plot of  $\log(-dc/dt)$  and  $\log [KBrO_3]$  in oxidation of gabapentin



**Fig. 2:** Plot of  $(-dc/dt)$  and  $[Ir(III)]$  in oxidation of gabapentin



**Fig. 3:** Plot of  $(-dc/dt)$  and  $[H^+]$  in oxidation of gabapentin

substrate, indicating zero order in substrate i.e. gabapentin (Table-1). The rate of reaction increases as the concentration of Iridium (III) chloride is increased, It was observed that values of  $(-dc/dt)$  were doubled when the concentration of iridium(III) was made two times, showing first order dependence on  $IrCl_3$  indicating first order of catalyst i.e.  $Ir(III)$  chloride (table-1). Kinetic result obtained on varying concentrations of chloride ions indicates fractional positive

order of chloride ion, which implies that rate of reaction increases when the concentration of  $Cl^-$  is increased (table-2). With increasing the concentration of  $[H^+]$ , the value of reaction rate also increases (Table-2). This showed first order dependence of  $[H^+]$  on the rate of oxidation of gabapentin. The rate measurements were taken at  $30^{\circ}$ - $45^{\circ}C$  and specific rate constants were used to draw a plot of  $\log k$  vs.  $1/T$  which was linear (Fig-2). The value of energy of Activation ( $\Delta E^*$ ) Arrhenius factor (A), entropy of activation ( $\Delta S^*$ ) and free energy of activation ( $\Delta G^*$ ) were calculated from rate measurement at  $30^{\circ}$ ,  $35^{\circ}$ ,  $40^{\circ}$ ,  $45^{\circ}C$  and these values have been recorded in Table-3. Moderate  $\Delta H^*$  and  $\Delta S^*$  values are favourable for electron transfer reaction. The value of  $\Delta H^*$  was due to energy of solution changes in the transition state. The high positive values of change in free energy of activation ( $\Delta G^*$ ) indicates highly solvated transition state, while fairly high negative values of change in entropy of activation ( $\Delta S^*$ ) suggest the formation of an activated complex with reduction in the degree of freedom of molecule. Negligible effect of mercuric acetate excludes the possibility of its involvement either as a catalyst or as an oxidant because it does not help the reaction proceed without potassium bromate. Hence the function of mercuric acetate is to act as a scavenger for any  $[Br^-]$  ion formed in the reaction. It helps to eliminate the parallel oxidation by  $Br_2$  which would have been formed as a result of interaction between  $Br^-$  and bromate ion. Potassium bromate has been used as an oxidant for a variety of compounds in acidic media (Natarajan and Venkatasubramaniam, 1969) sometimes in the presence of a catalyst. In alkaline and acidic medium, potassium bromate is ionised:



The  $BrO_3^-$  species has been reported to act as an oxidising agent in acidic (Anandan and Gopalan, 1985) as well as in alkaline medium (Singh and Srivastava, 1991).  $Ir(III)$  chloride has been reported to give a number of possible chloro species dependent on pH of the solution. Under the experimental pH range in the present investigation  $[Ir(Cl)_5(H_2O)]^{2-}$  has been proposed and confirmed as the reactive species dominant in the pH range 1.0 to 3.0. The kinetic results have been reported in Tables 1, 2 and 3.

### Mechanism

Considering the reaction steps shown in **scheme 2** and applying the steady state treatment, with reasonable approximation, the rate law may be written as shown in equations below.

$$Rate = k_3[C_2][HBrO_3] \quad (i)$$

$$[Ir(III)]_T = [C_1] + [C_2] \quad (ii)$$

$$\frac{d[C_1]}{dt} = k_{-1}[C_2] - k_1[C_1][Cl^-] \quad (iii)$$

$$[C_1] = \frac{k_{-1}[C_2]}{k_1[Cl^-]} \quad (iv)$$

$$[C_1] = \frac{[C_2]}{K_1[Cl^-]} \quad (v)$$

Where,  $K_1 = k_1/k_{-1}$

$$[Ir(III)]_T = [C_2] \left[ \frac{1 + K_1[Cl^-]}{K_1[Cl^-]} \right]$$

$$[C_2] = \frac{[Ir(III)]_T K_1 [Cl^-]}{1 + K_1 [Cl^-]}$$

$$\frac{d[HBrO_3]}{dt} = \frac{k_2[H^+][BrO_3^-]}{k_{-2}[HBrO_3]}$$

$$[HBrO_3] = K_2[H^+][BrO_3^-]$$

Putting the values of  $[C_2]$  and  $[HBrO_3]$  in equation (i), we get :

$$Rate = \frac{K_1 K_2 k_3 [Ir(III)]_T [Cl^-] [H^+] [BrO_3^-]}{1 + K_1 [Cl^-]}$$




Fig. 4: Plot of (a+bx) and  $[KBrO_3] \times 10^3$  M in oxidation of gabapentin




Fig. 5: Arrhenius plot for the oxidation of gabapentin

## CONCLUSION

In the present study of kinetic and thermodynamic properties pharmaceutical drug (gabapentin) by potassium bromate ( $KBrO_3$ ) in presence of Micro amount of Ir(III)

**Table 3:** Activation parameters for Ir(III) chloride catalyzed oxidation of gabapentin by  $KBrO_3$  at 35°C

Parameters	Temperature(T°C)	Gabapentine(-dc/dt)x 10 <sup>7</sup>
$k_1 \times 10^4 s^{-1}$	30 <sup>0</sup>	1.65
$k_1 \times 10^4 s^{-1}$	35 <sup>0</sup>	2.00
$k_1 \times 10^4 s^{-1}$	40 <sup>0</sup>	3.21
$k_1 \times 10^4 s^{-1}$	45 <sup>0</sup>	4.80
log A	...	12.02
$\Delta E^*$ (kJ mol <sup>-1</sup> )	35 <sup>0</sup>	69.13
$\Delta G^*$ (kJ mol <sup>-1</sup> )	35 <sup>0</sup>	74.83
$\Delta H^*$ (kJ mol <sup>-1</sup> )	35 <sup>0</sup>	73.47
$\Delta S^*$ (JK <sup>-1</sup> mol <sup>-1</sup> )	35 <sup>0</sup>	-18.392

chloride as catalyst in acidic medium. The following conclusions can be drawn:

1.  $[\text{IrCl}_6]^{3-}$  is considered as the reactive species of Ir(III) in acidic medium.
2.  $\text{HBrO}_3$  is the reactive species of potassium bromate in acidic medium.
3. The stoichiometry of the reaction was found to be 2:1 and the oxidation products of gabapentin were identified.
4. Activation parameters were computed from the Arrhenius plot.
5. The observed results have been explained by a plausible mechanism and the related rate law has been derived.

## REFERENCES

- Bai, W-S., Zhai, Y-Q., Liu, H-M., Lu H-Y., Song. W.Y., 2008. Kinetic and mechanism of Ir(III) catalysed oxidation of ethylene glycol by cerium(IV) in sulfuric acid media. *Indian Journal of Chemistry* 47 A 1194-1198.
- Singh, B., Srivastava, S., 1991. Kinetics and mechanism of ruthenium tetroxide catalysed oxidation of cyclic alcohols by bromate in a base. *Transition Metal Chemistry* 16, 466.
- Chaturvedi, A.K., Mishra, R. N., 2010. Kinetic and mechanism of Ir(III) chloride catalysed oxidation of ethyl amine by sodium meta periodate in acidic media. *Indian journal of Scientific Research* 1(2), 57-62.
- Chanakira, A., Chikwana, E., Peyton, D.H., Simoyi, R. H., 2006. Oxyhalogen-sulfur chemistry-Kinetics and mechanism of oxidation of methionine by aqueous iodine and acidified iodate. *Canadian journal of Chemistry*. 84, 49.
- Goel, A., Sharma, R., 2012. A kinetic and mechanistic study on the oxidation of arginine and lysine by hexa cyano ferrate(III) catalysed by iridium(III) in aqueous alkaline medium. *Journal of Chemical Engineering Materials Science* 3(1), 10.
- Hosahalli, R.V., Byadagi, K. S., Nandibewoor, S., Chimatadar, S.A., 2012. Kinetics and mechanism of ruthenium(III) catalyzed oxidation of Chloramphenicol-An antibiotic drug by diperiodato-cuprate(III) in aqueous alkaline medium. *Kinetic Catalysis* 53, 65.
- Jensen, A.A., Mosbacher, J., Elg, S., 2002. The Anti convulsant Gabapentin (Neurontin) Does Not Act through Y. Aminobutyric Acid-B Receptor. *Molecular. Pharmacology* 61, 1377.
- Koli, B.I., Nandibewoor, S.T., 2009. Kinetics and mechanism of Palladium(II) Catalysed Oxidation of dimethyl Sulfoxide by alkaline periodate. *Indian Journal of Chemistry* 48A 958.
- Kaushik, R.D., Kaur, M., Malik, R., Kumar, A., 2010. Kinetics and Mechanism of Mn(II) Catalyzed Periodate Oxidation of p-anisidine Effect of PH. *International Journal of Chemical Science* 8, 1379.
- Munavalli, D.S., Naik, P.N., Chimatadar, S. A., Nandibewoor, S. T., 2008. Kinetics and mechanistic study of iridium(III) Catalysed oxidation of sulphanilic acid by diperiodate cuprate(III) in aqueous alkaline medium *Reaction Kinetic Catalytic Letter* 94(2), 359.
- Mohana, K.N., Ramya, K.R., 2009. Ruthenium(III) Catalyzed oxidative cleavage of Thiamine hydrochloride with N-Bromo succinimide in presence of hydrochloride acid medium: A Kinetic and mechanistic. *Journal of Molecular Cataysis A: Chemical* 302, 80.
- Munikyamba, P., Rao, P.R., Sundaram, E.V., 1983. Kinetics of iodate oxidation of lactose in presence of  $[\text{IrCl}_3(\text{H}_2\text{O})_2\text{OH}]^-$  as homogeneous catalyst. *Journal of Indian Chemical Society* LX, 652.
- Munikyamba, P., Rao, P.R. Sundaram, E.V., 1981. Kinetics of iodate oxidation of lactose in presence of  $[\text{IrCl}_3(\text{H}_2\text{O})_2\text{OH}]^-$  as homogeneous catalyst. *Indian Journal of Chemistry* 20A, 1217.
- Mambo, E., Simoyi, R. H., 1993. Kinetics and Mechanism of the Complex Oxidation of Amino methane sulfinic Acid by Iodate in Acidic Medium. *Journal Physical Chemistry* 97, 3662.
- Muthakia, G.K., Jonnalagadda, S.B., 1989. Kinetics and mechanism of indigo Carmine-acidic reaction. An indicator Reaction for catalytic determination of Ru(III) ion. *International Journal of Chemical Kinetic* 21, 519.
- Wen-Yu, S., Rong-Hui, Z., Qing-Mei, J., 2005. Catalyzed Oxidation of Tetrahydro furfuryl Alcohol by Cerium(IV) in Aqueous Sulphuric Acid Medium. *Acta Physico-Chimica Sinica* 21 (08) 929.
- Singh, A., Rahmani. K., Singh, S., Singh, B., Singh, M., 2004. Mechanism of Ir(III)-catalysed and Hg(II)-co-catalysed oxidation of reducing sugars by N-bromoacetamide in acidic medium. *Journal of Physical Organic Chemistry* 17, 249.
- Singh, A., Singh. K., Srivastava, R., Rahmani, J., Srivastava, S., 2007. Mechanistic studies of oxidation of maltose and lactose by  $[\text{H}_2\text{OBr}]^+$  in presence of chloro-complex of Rh(III) as homogeneous catalyst *Journal of Organometallic Chemistry* 692, 4270.
- Singh, B., Singh, A.K., Singh, A., 2011. Mechanistic studies of aquachlororhodium(III) catalysis in potassium iodate oxidation of 1,2 propanediol in acidic medium: a kinetic approach *Indian Journal of Chemistry* 50A, 650.

Visit us at: <http://bosajournals.com/chemint/>  
 Submissions are accepted at: [editorci@bosajournals.com](mailto:editorci@bosajournals.com)